

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 18(3), Sept-Dec, 2020]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology, Pantnagar



ADVISORYBOARD

Patron

Dr. Tej Partap, Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Dr. A.S. Nain, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. A.K. Sharma, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. S.K. Kashyap, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. N.S. Jadon, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. K.P. Raverkar, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. Sandeep Arora, Ph.D., Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. Alaknanda Ashok, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. Alka Goel, Ph.D., Dean, College of Home Science, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. R.S. Chauhan, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India
Dr. R.S. Jadaun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

EDITORIALBOARD

Members

Prof. A.K. Misra, Ph.D., Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India
Dr. Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India
Dr. Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India
Dr. Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A
Dr. B.B. Singh, Ph.D., Visiting Professor and Senior Fellow, Dept. of Soil and Crop Sciences and Borlaug Institute for International Agriculture, Texas A&M University, U.S.A.
Prof. Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India
Dr. D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India
Dr. Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium
Dr. Desirazu N. Rao, Ph.D., Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India
Dr. G.K. Garg, Ph.D., Dean (Retired), College of Basic Sciences & Humanities, G.B. Pant University of Agric. & Tech., Pantnagar, India
Dr. Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines
Dr. Indu S Sawant, Ph.D., Director, ICAR - National Research Centre for Grapes, Pune, India
Dr. Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India
Dr. M.P. Pandey, Ph.D., Ex. Vice Chancellor, BAU, Ranchi & IGKV, Raipur and Director General, IAT, Allahabad, India
Dr. Martin Mortimer, Ph.D., Professor, The Centre of Excellence for Sustainable Food Systems, University of Liverpool, United Kingdom
Dr. Muneshwar Singh, Ph.D., Project Coordinator AICRP- LTFE, ICAR - Indian Institute of Soil Science, Bhopal, India
Prof. Omkar, Ph.D., Professor, Department of Zoology, University of Lucknow, India
Dr. P.C. Srivastav, Ph.D., Professor, Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India
Dr. Prashant Srivastava, Ph.D., Cooperative Research Centre for Contamination Assessment and Remediation of the Environment, University of South Australia, Australia
Dr. Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, U.S.A.
Dr. R.C. Chaudhary, Ph.D., Chairman, Participatory Rural Development Foundation, Gorakhpur, India
Dr. R.K. Singh, Ph.D., Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India
Prof. Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.
Dr. S.N. Maurya, Ph.D., Professor (Retired), Department of Gynecology & Obstetrics, G.B. Pant University of Agric. & Tech., Pantnagar, India
Dr. Sham S. Goyal, Ph.D., Professor (Retired), Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.
Prof. Umesh Varshney, Ph.D., Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India
Prof. V.D. Sharma, Ph.D., Dean Academics, SAI Group of Institutions, Dehradun, India
Dr. V.K. Singh, Ph.D., Head, Division of Agronomy, ICAR-Indian Agricultural Research Institute, New Delhi, India
Dr. Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological Agricultural Engineering, Texas A&M University, U.S.A.
Dr. Vinay Mehrotra, Ph.D., President, Vinlax Canada Inc., Canada

Editor-in-Chief

Dr. Manoranjan Dutta, Head Crop Improvement Division (Retd.), National Bureau of Plant Genetic Resources, New Delhi, India

Managing Editor

Dr. S.N. Tiwari, Ph.D., Professor, Department of Entomology, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Dr. Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

Dr. S.D. Samantray, Ph.D., Professor, Department of Computer Science and Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

CONTENTS

Marker assisted selection for aromatic and semi-dwarf segregants in cross of aromatic Katarni rice SUNDARAM BHARTI, P.K. SINGH, KUMARI SUVIDHA, SATYENDRA, S. P. SINGH, ANAND KUMAR and MANKESH KUMAR	188
D² and principal component analysis for variability studies in <i>Vigna</i> and <i>Phaseolus</i> species PRIYANKA BHARETI, R. K. PANWAR, ANJU ARORA and S. K. VERMA	193
Assessment of genetic parameters in F₅ recombinants derived from <i>Indica</i> rice (<i>Oryza sativa</i> L.) line Pusa 6A PRACHI PRIYA, MANKESH KUMAR, TIRTARTHA CHATTOPADHYAY, BISHUN DEO PRASAD, SWETA SINHA, ANAND KUMAR and SATYENDRA	198
Genetic diversity analysis by D² clustering of fodder yield and its related traits in forage sorghum HARSH DEEP, INDRANI CHAKRABORTY, SATYAWAN ARYA, PUMMY LAMBA, S. K. PAHUJA and JAYANTI TOKAS	203
Genetic diversity for morpho-physiological and seed vigour traits in wheat (<i>Triticum aestivum</i> L.) PUNEET KUMAR, Y.P.S. SOLANKI, VIKRAM SINGH and ASHISH	209
<i>In vitro</i> plant regeneration from mature embryo using different plant growth regulators in wheat genotype HD 3059 SWATI SHARMA, ASHWANI KUMAR, ANIL SIROHI, R. S. SENGAR, KAMAL KHILARI, MUKESH KUMAR and MANOJ K. YADAV	215
Weed management and crop geometry effect on nutrient uptake and yield in aerobic rice VASUNDHRA KAUSHIK, S. P. SINGH, V. P. SINGH, TEJ PRATAP and B. S. MAHAPATRA	222
Studies on sucker control in natu tobacco (<i>Nicotiana tabacum</i> L.) under rainfed vertisols S. JAFFAR BASHA, P. PULLI BAI, S. KASTURI KRISHNA and C. CHANDRASEKHARA RAO	228
Seed and oil yield of bidi tobacco (<i>Nicotiana tabacum</i> L.) varieties as influenced by planting geometry and fertilizer levels under rainfed vertisols S. JAFFAR BASHA, P. PULLI BAI, S. KASTURI KRISHNA and C. CHANDRASEKHARA RAO	232
Comparison of non-linear models on area, production and productivity of sugarcane crop in Uttar Pradesh JHADE SUNIL and ABHISHEK SINGH	237
Performance of improved varieties of true Cinnamon (<i>Cinnamomum verum</i> J. Presl.) in Andaman Islands, India AJIT ARUN WAMAN, POOJA BOHRA and R. KARTHIKA DEVI	243
Changing climate and its effect on rice yield in Meghalaya DEOTREPHY K. DKHAR, SHEIKH MOHAMMAD FEROZE, RAM SINGH and LALA I.P. RAY	249
Age related changes in morphometrical studies on ductus deferens of guinea fowl (<i>Numida meleagris</i>) TAMILSELVAN S, B. S. DHOTE and MEENA MRIGESH	257

Occurrence of gastrointestinal nematodes in goats slaughtered at Rewa, India D. MARAVI, A. K. DIXIT and POOJA DIXIT	261
Autoimmune haemolytic anaemia in a dog-A case report NEERAJ KUMAR, MUNISH BATRA and R.S. CHAUHAN	265
Erythrocytic anaplasmosis with <i>Fasciolosis</i> in a cross-bred cattle: A case report NEERAJ KUMAR and MUNISH BATRA	269
Modification and evaluation of Pant-ICAR controlled traffic seed-cum-deep fertilizer applicator for multi-crop seeder-cum-deep placement of fertilizers applicator MANISH KUMAR, T.C THAKUR, MANOJ KUMAR and SATYA PRAKASH KUMAR	272
Drying characteristics of shrimp (<i>Metapenaeus dobsoni</i>) in electrical dryer D.S. ANIESRANI DELFIYA, S. MURALI, P.V. ALFIYA and MANOJ P. SAMUEL	281
Baur dam breach analysis using various Manning's roughness values MEENAKSHI RAMOLA, JYOTHI PRASAD and H. J. SHIVA PRASAD	286
Study of constipation and related factors among female students of Pantnagar RITA SINGH RAGHUVANSHI, NIDHI JOSHI, DIKSHA SINGH, SHIKHA SINGH, MEENAL and DASHRATH BHATI	290
Work -related musculoskeletal disorders among chikankari workers in Lucknow (U.P.) POONAM SINGH and KATYAYNI	297
Technology adoption and productivity enhancement in groundnut cultivation: An impact assessment of farm women groups K.UMA, T. NIVETHA and S. PRAVEENA	302
Health hazard and constraints of chikankari worker in Lucknow (U.P.) POONAM SINGH and KATYAYNI	310
Studies on Indigenous Agricultural Technical Knowledge prevalent among the farmers of Assam for the management of common pests and diseases in major crops DEVAMITRA TARAFDAR and NIRMAL MAZUMDER	315
Television viewing pattern among students of CCS Haryana Agricultural University, Hisar ANIL KUMAR MALIK, KRISHAN YADAV and SUNIL KUMAR	325
Media content development and it's standardization for farmers REETA DEVI YADAV, GEETAMATI DEVI and RITA GOAL	331
Analysis of learning behavior and pattern of online learners on a MOOC platform G.R.K. MURTHY, SEEMA KUJUR, S. SENTHIL VINAYAGAM, YASHAVANTH B.S., CH. SRINIVASA RAO, P. S. PANDEY, VANITA JAIN and INDRADEVI T.	338

Changing climate and its effect on rice yield in Meghalaya

DEOTREPHY K. DKHAR², SHEIKH MOHAMMAD FEROZE¹, RAM SINGH³ and LALA I.P. RAY⁴

¹College of Agriculture, ^{2,3,4}School of Social Sciences, College of Post Graduate Studies, Central Agricultural University – Imphal, Umiam-793103 (Meghalaya)

ABSTRACT: The change in precipitation pattern and temperature directly and indirectly affect the weather dependent agriculture sector. So, this study assessed the effect of rainfall and temperature variability on the mean and variance of rice yield in Meghalaya by using Just & Pope model. The state received fairly good amount of rainfall during rice growing season but the minimum and maximum temperatures have increased during 1975-2007. Average monsoon maximum temperature has positive and significant effect on the mean yield of *kharif* rice. The significant effect of the squared term of monsoon minimum temperature, and the interaction term of monsoon minimum and monsoon maximum temperature implies their non-linear effect on the mean yield of *kharif* rice. The variance function reveals that the climatic parameters along with time trend are risk decreasing factors but only time trend is statistically significant.

Key words: Just and Pope, Meghalaya, rainfall, Sen's slope, temperature

The crop yield is a function of different inputs *viz.*, seed, nutrients, water, labour etc. Precipitation and temperature are the two important weather parameters which have direct bearing on agricultural production (Deschenes and Greenstone, 2006). The studies on effect of climatic parameters on mean crop yield has taken centre stage in the regime of climate change and is important for rainfed agriculture. Changes in precipitation and temperature have direct effect on crop water use and impact the crop yield (Birthal *et al.*, 2014; Grover and Upadhyay, 2014). Rice yields are projected to decline by 5-12 per cent over India and China (Lin *et al.*, 2004) and by 4 per cent in overall Asia by the end of the century (Murdiyarso, 2000). Saito *et al.* (2006) reported that rice yield is closely associated with the total amount of rainfall from June to August in northern Laos, since this period coincides with the rapid vegetative and early reproductive growth stages. They reported that if total rainfall from June to August was less than 610 mm the average yield of rice was 1.4 MT/ha, whereas if rainfall was greater than 690 mm the average yield increased to 2.5 MT/ha. Birthal *et al.* (2014) has studied the effect of temperature on crop yield in India during the period of 1969-2005 and found that rise in maximum temperature has a negative and significant effect on yields of *kharif* as well as *rabi* crops. On the other hand, a rise in minimum temperature has a significantly positive impact on yields of most crops. The opposing effects of rise in minimum and maximum temperatures suggest that temperature has a non-linear effect on the crop yields. Kumar *et al.* (2014) estimated that in India as a result of a 10 per cent deficit in rainfall during 1980 to 2005, the yield of rice has declined by 6.3 per cent, and

also reported that 10 per cent drought intensity would be responsible for the fall in production of rice by 10 per cent.

Meghalaya is an agrarian state in the north-eastern hill (NEH) region of India. The contribution of agriculture in state GDP is 18.06 per cent (GoI, 2015) and about 60 per cent of the population is dependent on agriculture for livelihood. Rice is the staple food for the people and the primary cereal crop cultivated in the state with 108.27 thousand ha area under rice. The crop is cultivated primarily in rain-fed condition as the area under irrigation is only 22.66 per cent as in 2011. Alike other states of the region Meghalaya also faces the challenge in terms of change in climatic parameters; specifically precipitation and temperature (Jain *et al.*, 2013; Chakraborty *et al.*, 2017). Rice being susceptible to climate change and events like droughts and floods (Karim *et al.*, 1996; Yu *et al.*, 2010; Sarker *et al.*, 2014); any change in climatic parameters is expected to have negative effect on rice yield and in turn the household food availability. Hence, in this paper we have studied the trend in climatic parameters and their effect not only in mean yield but also on variance of yield of rice in Meghalaya. This kind of empirical study is lacking for the state.

MATERIALS AND METHODS

Data

Meghalaya situated in the North Eastern Indian Himalaya is one of the seven sister hill states which lies between

24°57'N to 26°10'N latitude and 89°46'E to 92°52'E longitudes. The elevation of the plateau ranges between 150 m to 1961 m above mean sea level (msl). The state receives very high rainfall and known as the “abode of clouds” aptly. Secondary data on daily rainfall (1975-2007) and daily temperature (1975-2009) were extracted from high resolution $1^\circ \times 1^\circ$ daily gridded data obtained from India Meteorological Department (IMD), Pune for three stations i.e., 90.5°E longitude and 25.5°N latitude, 91.5°E longitude and 25.5°N latitude and 92.5°E longitude and 25.5°N latitude. The individual station data were used to arrive at state average. The secondary data on yield of rice was collected from the Directorate of Economics and Statistics, Government of Meghalaya to estimate the impact of rainfall on rice yield.

Normality Test

Normal Quantile-Quantile (QQ) plot: It's a visual check in which a scatter diagram of two sets of quantiles i.e., the theoretical or normal quantile and sample quantile are plotted against each other. Hence if the distribution is normal the sample quantiles will be around the normal quantile i.e., the diagonal straight line.

JarqueBera (JB) test of normality test: At first the residuals are calculated using ordinary least squares (OLS); and then skewness and kurtosis is calculated. The test statistics is as follows:

$$JB = n \left[\frac{S^2}{6} + \frac{(K-3)^2}{24} \right]$$

Where n = sample size, S = skewness coefficient, K = kurtosis coefficient. The null hypothesis is the residuals are normally distributed. In that case $S=0$, $K=0$ and $JB=0$. (Gujarati, 2003)

Trend analysis

The parametric tests are robust than non-parametric tests but non-parametric tests are useful when the variables do not follow normal distribution. Hence, we have calculated trends for yield and all the climatic variables using linear regression equations (parametric) as well as monotonic trends using Sen's slope (non-parametric).

Linear regression equation: The simple linear regression model used is as follows:

$$Y_i = \beta_1 + \beta_2 \text{ Time} \dots\dots\dots(2)$$

Where Y_i is regress and, β_1 is the intercept and β_2 is the slope coefficient. The parameters are estimated using ordinary least squares (OLS) technique and tested for their significance using corresponding probability (p) values.

Sen's slope and Man-Kendall test

Sen's slope estimates the magnitude of trend (Sen, 1968) which is used by a number of researchers in hydro-meteorological studies (Yue and Hashino, 2007; Partal and Kahya, 2006; Jhalaria and Singh, 2010; Jain and Kumar, 2012; Chakraborty *et al.*, 2014). In this method at first a set of monotonic slope is calculated as

$$T_i = \frac{X_j - X_k}{j - k} \text{ for } i = 1, \dots, N \dots\dots(3)$$

Where, x_j and x_k are data values at time j and k ($j > k$), respectively. The median of these N values of T_i is Sen's estimator of slope, which is calculated as follows:

$$\beta = \begin{cases} \frac{T_{N+1}}{2} & ; N \text{ is odd} \\ \frac{1}{2 \left(\frac{T_N}{2} + \frac{T_{N+1}}{2} \right)} & ; N \text{ is even} \dots\dots\dots(4) \end{cases}$$

The significance of Sen's slope estimates are tested using Mann Kendal (MK) Z statistic (Mann, 1945; Kendall, 1970). The statistics (S) is defined as (Salas, 1993) follows:

$$S = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \text{sign}(x_j - x_i) \dots\dots\dots(5)$$

Where, N is the number of data points. Assuming $(x_j - x_i) = \theta$, the value of sign (θ) is computed as follows:

$$\text{sign}(\theta) = \begin{cases} 1 & \text{if } \theta > 0 \\ 0 & \text{if } \theta = 0 \\ -1 & \text{if } \theta < 0 \end{cases} \dots\dots\dots(6)$$

This statistics represents the number of positive differences minus the number of negative differences for all the differences considered.

The standard normal deviate (Z-statistics) is computed as follows:

$$Z = \begin{cases} \frac{s-1}{\sqrt{\text{var}(s)}} & \text{if } S > 0 \\ 0 & \text{if } S = 0 \\ \frac{s+1}{\sqrt{\text{var}(s)}} & \text{if } S < 0 \end{cases} \dots\dots\dots(7)$$

Where, $\text{var}(s) = \frac{1}{18} [N(N-1)(2N+5) - \sum_{i=1}^n t_i(t_i-1)(2t_i+5)]$

If the computed value of $|Z| > z_{\alpha/2}$, the null hypothesis (H_0) is rejected at α level of significance in a two-sided test.

Effect of climatic variability on rice yield: Just-Pope model

Just-Pope stochastic production function was employed to estimate the effect of climatic variability not only on the mean yield but also yield variability by many researchers across the globe (Tveteras, 1999; Chen *et al.*, 2004; Isik and Devadoss, 2006; Kim and Pang, 2009). The Cobb–Douglas and linear–quadratic forms are consistent with the Just and Pope postulates (Tveteras, 2000; Kim and Pang, 2009). The function is as follows

$$y = f(X; \beta) + \omega h(X; \delta)^{0.5} \dots\dots (i)$$

Where,

y is the productivity (production per hectare) of *kharif* rice, X is a vector of explanatory variables ω is the stochastic term with mean zero and variance 1, β and W are the production function parameters to be estimated using historical data (Just and Pope, 1978). From equation (i), the expected productivity of crop is given by $E(y) = f(X)$ and crop variability is given by $V(y)=h(X;\beta)$. The derivatives of the variance function $h(X;\beta)$ w.r.t. the input variables, viz., precipitation and temperature can be used to identify whether a climate variable increases or decreases crop variability. So if, $h_x = \delta h/\delta x > 0$, it indicates that the corresponding input variable x is risk increasing, if $h_x < 0$ it implies risk decreasing.

As the equation (i) has heteroskedastic errors; Feasible Generalized Least Squares (FGLS) technique has been applied to estimate the mean and variance functions and the model was estimated using GRETL 1.10.1. The variables used in the Just and Pope model and their functional forms are as below:

RESULTS AND DISCUSSION

Descriptive statistics of the variables

Table 1: Descriptive statistics for yield and climatic variables

Variables	Average	Extreme value		CV (%)
		Minimum	Maximum	
Yield (MT/ha)	1.34	0.90	1.96	24.01
MRain (mm)	1614.78	957.80	3330.93	29.80
MinT (°C)	23.37	22.77	24.69	0.43
MaxT (°C)	30.44	29.82	31.64	1.32

& * represents $p<0.05$ and $p<0.01$, respectively




Fig 1: Trend in *kharif* rice yield in Meghalaya

The *kharif* rice yield has increased from 1.13 MT/ha in 1975 to 1.83 MT/ha in 2007 (Fig. 1). The estimated annual trend growth of yield, calculated using LOGEST function in excel, is 1.73 per cent. This increase can be attributed to technological interventions in terms of high yielding varieties, irrigation and fertilizer application. The highest yield of 1.96 MT/ha was registered in 2005 which was a high rainfall year whereas, the lowest yield of 1.73 MT/ha in 1986. The instability in yield was very high during 1975-2007 which can be observed in Fig. 1 and is evident from the high value of estimated coefficient of variation (CV) for the yield (Table 1). Similarly, the inter year variations in monsoon rainfall was very high (CV=30 %) during the study period. The average monsoon rainfall was 1615 mm which is about 69 per cent of the total annual rainfall in Meghalaya during the period 1975-2007. The state received highest monsoon rainfall of 3331mm in 2005 and it was followed by minimum monsoon rainfall of 958 mm in the very next year i.e., 2006. The average MinT and MaxT were 23.37°C and 30.44°C, respectively during 1975-2007 (Table 1). The inter year variation in MaxT was relatively higher than MinT in the state.

Trend in variables (Fig 2 to Fig 4)




Fig 2: Trend in monsoon rainfall in Meghalaya

The parametric trends in Yield, MRain, MinT and MaxT are presented in Fig. 1 to Fig.4 which depicts positive linear trends for all these variables. The normality of the variables is checked first visualizing the Normal Q-Q plot and further confirming it by applying JarqueBera (JB) test of normality. The Normal Q-Q plot reveals that the residuals are around the diagonal line for all the variables

Variables	Functional form	
	Mean function	Variance function
Dependent : <i>Kharif</i> yield (1975-2007)		
Explanatory		
Time	Time, MRain, MaxT, MinT, squared MaxT,	Time, MRain, MaxT, MinT
Monsoon rainfall (MRain) in mm	squared MinT, squared MRain, MaxT*MinT,	
Monsoon maximum temperature (MaxT) in °C	MaxT*MRain, MinT*MRain.	
Monsoon minimum temperature (MinT) in °C		




Fig 3: Trend in monsoon minimum temperature in Meghalaya

except MaxT. Similarly, the JB statistics are also significant for all the variables except MaxT. Hence, it is concluded that only MaxT follows normal distribution (Fig. 5 and Table 2).




Fig 4: Trend in monsoon maximum temperature in Meghalaya

The positive monotonic trend is significant for yield ($p < 0.05$) and the yearly increment is 0.03MT/ha. Though the MRain shows increasing linear monotonic trend but it is statistically insignificant. Similar finding was reported by Das *et al.* (2015) who observed positive but insignificant trend in monsoon rainfalls during 1960-2010 at Cherrapunjee and Shillong in Meghalaya. Chakraborty *et al.* (2017) reported negative trend in monsoon rainfall for Umiyam in Meghalaya is insignificant, which is similar to the findings of Jain *et al.* (2013) for Assam & Meghalaya sub-division for the period of 1871-2008.

The MinT and MaxT have shown significant positive monotonic ($p < 0.05$) and positive linear ($p < 0.05$) trend. The average monsoon minimum and maximum temperature have increased by 0.3°C/decade and 0.02°C/decade, respectively (Table 2 and Fig. 4). Chakraborty *et*

al. (2014) analysed the monthly, seasonal as well as the annual trends in mean air temperature over seven different places situated at various hills of northeastern India. They reported a dissimilar pattern of changes among the stations; places viz., Basar in Arunachal Pradesh, Imphal in Manipur and Gangtok in Sikkim where the mean temperature was lower, i.e., those places which are climatologically cooler, the increase in temperature is significant. Jain *et al.* (2013) also observed increasing trend in temperature in Assam & Meghalaya subdivision. Similarly, Jhaharia and Singh (2010) reported increase temperature trend in Assam which is different in physiographic.

Relationship between yield and climatic variables

Table 3: Zero order correlation coefficients among the variables

Variables	Yield	MRain	MinT	MaxT
Yield	1.00			
MRain	0.22	1.00		
MinT	0.62***	0.44**	1.00	
MaxT	0.51**	0.20	0.61***	1.00
	dYield	MRain	d_MinT	d_MaxT
d_Yield	1.00			
MRain	0.31	1.00		
d_MinT	0.18	-0.04	1.00	
d_MaxT	0.02	0.21	0.23	1.00

& * represents $p < 0.05$ and $p < 0.01$, respectively

Note d_Yield, d_MinT and d_MaxT are the first difference forms

Table 4: Stationarity of different the variables: ADF test

Variables	Model	tau	P
Yield	with constant	-0.65	0.85
	with constant and trend	-1.75	0.71
MRain	with constant	-6.42	<0.01
	with constant and trend	-6.81	<0.01
MinT	with constant	0.84	0.99
	with constant and trend	-1.14	0.92
MaxT	with constant	-3.65	0.01
	with constant and trend	-4.06	0.02

The zero order correlation coefficients among the variables were calculated and presented in Table 3. The MinT and

Table 2: Normality test and estimated trend coefficients of the yield and climatic variables

Variables	Normality test		Parametric (linear)	Non-parametric (monotonic)	
	Jarque Berra (B)	P value	slope coefficient	Men Kendall's Z statistic	Sen's slope estimate
Yield (MT/ha)	4.32**	0.01	0.03***	2.93**	0.03
MRain (mm)	43.23***	<0.01	13.57	1.84	0.04
MinT (°C)	8.18**	0.02	0.03 ***	2.92**	0.03
MaxT (°C)	8.25	0.12	0.02 **	1.38	0.05

& * represents $p < 0.05$ and $p < 0.01$, respectively

MaxT are positively correlated ($p < 0.01$). The years with higher minimum temperature received significant ($p < 0.01$) higher monsoon rainfall in the state. The MaxT ($p < 0.05$) and MinT ($p < 0.01$) are significantly correlated with *kharif* rice yield (Table 3). Then, the stationarity for all the variables was checked using correlogram (see Annex I) and further applying Augmented Dickey-Fuller (ADF) test. It is found that all the variables, except yield of rice, were non-stationary in nature (Table 4). Hence, they were made stationary by taking the first difference of each series. But when the simple correlation coefficients were calculated using the stationary variables it revealed no significant linear association among them (Table 3).

Critical temperature for rice plant at different growth stages

Stage	Critical Temperature ⁰ C		
	Low	High	Optimum
Germination	16-19	45	18-40
Seedling emergence and establishment	12-35	35	25-30
Rooting	16	35	25-28
Leaf elongation	7-12	45	31
Tillering	9-16	33	25-31
Initiation of panicle primordia	15	-	-
Panicle differentiation	15-20	30	-
Anthesis	22	35-36	30-33
Ripening	12-18	>30	20-29

(Source: Yoshida 1978)

Effect on mean and variance in rice yield

Table 5: Estimated coefficients of Just and Pope model

Variables	Coefficient	p-value
Mean function		
Constant	-0.30	0.640
Time	<-0.01	0.882
MaxT	0.24**	0.050
MinT	0.12	0.717
MRain	<0.01	0.708
Squared MaxT	0.51	0.165
Squared MinT	0.15**	0.043
Squared MRain	<-0.01	0.836
MaxT*MinT	-0.99**	0.042
MaxT*MRain	0.99	0.199
MinT*MRain	<-0.01	0.178
Variance function		
Constant	-3.52	0.076
MRain	<-0.01	0.54
MaxT	-0.34	0.802
MinT	-0.64	0.616
Time	-0.03**	0.032
F value (10,21)	12.48***	<-0.01
Akaike criterion	186.06	

** & *** represents $p < 0.05$ and $p < 0.01$, respectively

The variables of Just and Pope Model are assumed to be stationary (Chen *et al.*, 2004) as non-stationary series may





Fig 5: Q-Q Plot for Yield, MRain, Min T and Max T

Correlogram for yield, MRain, MinT and MaxT



lead to spurious results (Chen and Chang, 2005). Hence, we used the original series of MinT which is stationary and first differenced form of Yield, MRain and MaxT. The coefficient of Just and Pope Model reveals that maximum temperature has a positive and significant effect on the mean yield of *kharif* rice in Meghalaya (Table 5). The significant effect of the squared term of monsoon minimum temperature, and the interaction term of monsoon minimum and monsoon maximum temperature implies their non-linear effect on the mean yield of *kharif* rice. The optimum temperature for rice cultivation is about 25-33°C though it varies across different stages (Yoshida, 1978) (see Annex II). The long term average temperature was 26.91°C in Meghalaya during the study period; hence the increasing temperature has positive effect on mean yield. The variance function too reveals that the climatic parameters along with time trend are risk decreasing factors but only time trend is statistically significant ($p < 0.05$). This means technological progress had played positive role in reducing the variance in yield of rice. This finding is in line of the finding of Sarker *et al.* (2014) who found that mean minimum temperature is a risk decreasing factor for *Aus* and *Aman* rice in Bangladesh.

CONCLUSION

The monsoon rainfall in Meghalaya showed increasing but insignificant trend during the period of 1975-2007. The linear trends for minimum and maximum temperatures are positive and significant. Monsoon maximum temperature has positive and significant effect on the mean yield of *kharif* rice. Time is risk decreasing factor for rice yield in Meghalaya. Hence, the study does not find significant negative effect of climatic variables on rice yield in Meghalaya as the average rainfall is sufficient and though the temperature is increasing but not crossed the critical limit to become detrimental to rice yield in the state.

ACKNOWLEDGEMENTS

The fellowship to first author from Central Agricultural University for MSc dissertation is acknowledged.

REFERENCES

- Birthal, P.S., Khan, M.T., Negi, D.S. and Aggarwal, S. (2014). Impact of climate change on yields of major food crops in India: Implications for food security. *Agricultural Economics Research Review*, 27: 145-155.
- Chakraborty, D., Saha, S., Singh, R.K., Sethy, B.K., Kumar, A., Saikia, U.S., Das, S.K., Makdoh, B., Borah, T.R., Chanu, N.A., Walling, I., Rollinganal, P.S., Chowdhury, S., and Daschoudhury, D. (2017). Spatio temporal trends and change point detection in rainfall in different parts of North Eastern Indian States. *Journal of Agrometeorology*, 19:160-163.
- Chakraborty, D., Singh, R.K., Roy, A., Saha, S., Sethy, B.K., Kumar, A. and Ngachan, S.V. (2014). Increase in extreme day temperature in hills of Meghalaya: Its possible ecological and biometeorological effect. *Journal of Agrometeorology*, 16:147-152.
- Chen, C., McCarl, B.A. and Schimmelpfennig, D.E. (2004). Yield variability as influenced by climate: A statistical investigation. *Climate Change*, 66: 239-261.
- Chen, C. and Chang, C. (2005). The impact of weather on crop yield distribution in Taiwan: Some new evidence from panel data models and implications for crop insurance. *Agricultural Economics*, 33: 503-511.
- Das, S., Tomar, C.S., Saha, D., Shaw, S.O. and Singh, C. (2015). Trends in Rainfall Patterns over North-East India during 1961-2010. *International Journal of Earth and Atmospheric Science*, 2: 37-48.
- Deschenes, O. and Greenstone, M. (2006). The economic impacts of climate change: Evidence from agricultural output and random fluctuations in weather. *American Economic Review*, 97: 354-385.
- GoI. (2015). Basic Statistics of North Eastern Region 2015. Issue No 14, Government of India,
- Grover, D.K. and Upadhyay, D. (2014). Changing climate pattern and its impact on paddy productivity In Ludhiana districts of Punjab. *Indian Journal of Agricultural Economics*, 69: 150-162.
- Gujarati, D.N. (2003). Basic Econometrics. 4th Edition, McGraw Hill, New York.
- Isik, M. and Devadoss, S. (2006). An analysis of the impact of climate change on crop yields and yield variability. *Applied Economics*, 38: 835-844.
- Jain, S.K. and Kumar, V. (2012). Trend analysis of rainfall and temperature data for India. *Current Science*, 102: 37-49.
- Jain, S.K., Kumar, V. and Saharia, M. (2013). Analysis of rainfall and temperature trends in northeast India. *International Journal of Climatology*, 33: 968-978. doi10.1002/joc.3483
- Jhajharia, D. and Singh, V.P. (2010). Trends in temperature, diurnal temperature range and sunshine duration in Northeast India. *International Journal of Climatology*, 31:1353-136.
- Just, R.E. and Pope, R.D. (1978). Stochastic representation

- of production functions and econometric implications. *Journal of Economics*, 7: 67-86.
- Karim, Z., Hussain, S.G. and Ahmed, M. (1996). Assessing impact of climate variation on food grain production in Bangladesh. *Journal of Water Air Soil Pollution*, 92:53–62.
- Kendall, M.G (1970). Rank Correlation Methods. 2nd Ed., New York: Hafner.
- Kumar, P., Joshi, P.K. and Aggarwal, P. (2014). Projected effect of drought on supply, demand, and prices of crops in India. *Economic and Political Weekly*, 49:55-63.
- Lin, E.D., Xu, Y.L., Ju, H. and Xiong, W. (2004). Possible adaptation decisions from investigating the impacts of future climate change on food and water supply in China. Proc of the AIACC Asia-Pacific region workshop, Nov 5-12, 2004. Manila, Philippines.
- Mann, H.B. (1945). Nonparametric tests against trend. *Econometrica*, 13:245:259.
- Murdiyarso, D. (2000). Adaptation to climate variability and change: Asian perspectives on agriculture and food security. *Environment, Monitoring and Assess*, 61:123-131.
- Partal, T. and Kahya, E. (2006). Trend analysis in Turkish precipitation data. *Hydrological Processes*, 20: 2011–2026.
- Kim, M. and Pang, A. (2009). Climate change impact on rice yield and production risk. *Journal of Rural Development*, 32:17–29.
- Sarker, M.A.R., Khorshed, A. and Gow.J. (2014). Assessing the effects of climate change on rice yields: An econometric investigation using Bangladeshi panel data. *Economic Analysis and Policy*, 44:405–416.
- Saito, K., Linqvist, B., Keobualapha, B., Phanthaboon, K., Shiraiwa, T. and Horie, T. (2006). Cropping intensity and rainfall effects on upland rice yields in northern Laos. *Plant Soil*, 284:175-185.
- Salas, J.D. (1993). Analysis and modeling of hydrologic time series. In: Maidment DR (ed), *Handbook of Hydrology*, McGraw-Hill, New York, 1993, Pp. 19.1–19.72.
- Sen, P.K. (1968). Estimates of the regression coefficient based on Kendall's tau. *Journal of American Statistical Association*, 39:1379–1389.
- Tveteras, R. (1999). Production risk and productivity growth: Some findings for Norwegian Salmon aquaculture. *Journal of Productivity Analysis*, 12:161–179.
- Tveteras, R. (2000). Flexible panel data models for risky production technologies with an application to Salmon aquaculture. *Econometrics Review*, 19: 367–389.
- Yoshida, S. (1978). Tropical climate and its influence on rice. IRRI Research Paper Series 20. Los Baños, Philippines, IRRI.
- Yue, S. and Hashino, M. (2007). Long term trends of annual and monthly precipitation in Japan, *Journal of the American Water Resource Association*, 39,587–596 doi 10.1111/j.1752-1688.2003.tb03677.x.
- Yu, W.H., Alam, M., Hassan, A., Khan, A.S., Ruane, A.C., Rosenzweig, C., Major, D.C. and Thurlow, J. (2010). Climate Change Risk and Food Security in Bangladesh. EarthScan, London.

Received: May 20, 2020

Accepted: November 21, 2020