

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 22(1) January-April 2024]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology, Pantnagar



ADVISORYBOARD

Patron

Dr. Manmohan Singh Chauhan, Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Dr. A.S. Nain, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. S.K. Kashyap, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. S.P. Singh, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. K.P. Raverkar, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. Sandeep Arora, Ph.D., Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. Alaknanda Ashok, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Dr. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

EDITORIALBOARD

Members

Prof. A.K. Misra, Ph.D., Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India

Dr. Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Dr. Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Dr. Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A

Dr. B.B. Singh, Ph.D., Visiting Professor and Senior Fellow, Dept. of Soil and Crop Sciences and Borlaug Institute for International Agriculture, Texas A&M University, U.S.A.

Prof. Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

Dr. D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Dr. Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Dr. Desirazu N. Rao, Ph.D., Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

Dr. G.K. Garg, Ph.D., Dean (Retired), College of Basic Sciences & Humanities, G.B. Pant University of Agric. & Tech., Pantnagar, India

Dr. Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Dr. Indu S Sawant, Ph.D., Director, ICAR - National Research Centre for Grapes, Pune, India

Dr. Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

Dr. M.P. Pandey, Ph.D., Ex. Vice Chancellor, BAU, Ranchi & IGKV, Raipur and Director General, IAT, Allahabad, India

Dr. Martin Mortimer, Ph.D., Professor, The Centre of Excellence for Sustainable Food Systems, University of Liverpool, United Kingdom

Dr. Muneshwar Singh, Ph.D., Project Coordinator AICRP- LTFE, ICAR - Indian Institute of Soil Science, Bhopal, India

Prof. Omkar, Ph.D., Professor, Department of Zoology, University of Lucknow, India

Dr. P.C. Srivastav, Ph.D., Professor, Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Dr. Prashant Srivastava, Ph.D., Cooperative Research Centre for Contamination Assessment and Remediation of the Environment, University of South Australia, Australia

Dr. Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, U.S.A.

Dr. R.C. Chaudhary, Ph.D., Chairman, Participatory Rural Development Foundation, Gorakhpur, India

Dr. R.K. Singh, Ph.D., Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Prof. Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

Dr. S.N. Maurya, Ph.D., Professor (Retired), Department of Gynecology & Obstetrics, G.B. Pant University of Agric. & Tech., Pantnagar, India

Dr. Sham S. Goyal, Ph.D., Professor (Retired), Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Prof. Umesh Varshney, Ph.D., Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

Prof. V.D. Sharma, Ph.D., Dean Academics, SAI Group of Institutions, Dehradun, India

Dr. V.K. Singh, Ph.D., Head, Division of Agronomy, ICAR-Indian Agricultural Research Institute, New Delhi, India

Dr. Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological Agricultural Engineering, Texas A&M University, U.S.A.

Dr. Vinay Mehrotra, Ph.D., President, Vinlax Canada Inc., Canada

Editor-in-Chief

Dr. Manoranjan Dutta, Head Crop Improvement Division (Retd.), National Bureau of Plant Genetic Resources, New Delhi, India

Managing Editor

Dr. S.N. Tiwari, Ph.D., Professor, Department of Entomology, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Dr. Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

Dr. S.D. Samantray, Ph.D., Professor, Department of Computer Science and Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

CONTENTS

Productivity, nutrient uptake and economics of sweet corn (<i>Zea mays</i> L. var. <i>saccharata</i>) under different planting geometry and NPK levels	1-7
AMIT BHATNAGAR, SAILESH DEB KARJEE, GURVINDER SINGH and DINESH KUMAR SINGH	
Integrated effect of natural farming concoctions and organic farming practices with various NPK doses on quality of bread wheat	8-13
PRERNA NEGI, MOINUDDIN CHISTI and HIMANSHU VERMA	
Characterization and fertility capability classification of some soils in the rain forest zone of Edo state, Nigeria	14-25
OKUNSEBOR, F.E., OGBEMUDIA, I. and OKOLIE, S. I.	
Characterization and classification of guava growing soils of North-East Haryana according to frame work of land evaluation (FAO, 1993)	26-35
DHARAM PAL, MANOJ SHARMA, R.S. GARHWAL and DINESH	
Interactive impact of heavy metals and mycorrhizal fungi on growth and yield of pepper (<i>Capsicum annuum</i> Linn.)	36-47
SHARMILA CHAUHAN, MOHINDER SINGH, SNEHA DOBHAL, DEEKSHA SEMWAL and PRAVEEN	
Response of chilli (<i>Capsicum annuum</i> var. <i>annuum</i> L.) to different nutrient management practices	48-58
SHEETAL, K.C. SHARMA, SHIVAM SHARMA, NEHA SHARMA, D.R. CHAUDHARY, SANDEEP MANUJA and AKHILESH SHARMA	
Trend detection in weather parameters using Mann-Kendall test for Tarai region of Uttarakhand	59-67
SHUBHIKA GOEL and R.K. SINGH	
Comparative study of antioxidant potential of fresh peel from different citrus species	68-74
TARU NEGI, ANIL KUMAR, ARCHANA GANGWAR, SATISH KUMAR SHARMA, ANURADHA DUTTA, NAVIN CHAND SHAHI, OM PRAKASH and ASHUTOSH DUBEY	
Suitability of Quinoa Grains (<i>Chenopodium Quinoa</i> Willd.) for development of Low Glycemic Index Biscuits	75-84
RUSHDA ANAM MALIK, SARITA SRIVASTAVA and MEENAL	
A study on dietary intake among school-going adolescent girls of Udaipur, Rajasthan during COVID-19	85-92
JYOTI SINGH and NIKITA WADHAWAN	
Nutritional and sensory evaluation of gluten free chapatti developed using underutilised food sources	93-98
AYUSHI JOSHI, ARCHANA KUSHWAHA, ANURADHA DUTTA, ANIL KUMAR and NAVIN CHANDRA SHAHI	
Nutrient-enriched wheat chapatti with fresh pea shells (<i>Pisum sativum</i> L.): A comprehensive quality assessment	99-109
AMITA BENIWAL, SAVITA, VEENU SANGWAN and DARSHAN PUNIA	

Pearl Millet-Based Pasta and Noodles Incorporated with <i>Jamun</i> Seed Powder: Quality Analysis SAVITA, AMITA BENIWAL, VEENU SANGWAN and ASHA KAWATRA	110-121
Unlocking the biofortification potential of <i>Serratia marcescens</i> for enhanced zinc and iron content in wheat grains BHARTI KUKRETI and AJAY VEER SINGH	122-131
Antioxidant and anti-inflammatory properties of sun-dried leaves and fruits of wild <i>Pyracantha crenulata</i> (D. Don) M. Roem. SUGANDHA PANT, PREETI CHATURVEDI, AAKANSHA VERMA, MANDEEP RAWAT, VAISHNAVI RAJWAR and KAVITA NEGI	132-141
Studies on productive herd life, longevity, and selective value and their components in crossbred cattle SHASHIKANT, C.V. SINGH and R.S. BARWAL	142-150
Studies on replacement rate and its components in crossbred cattle SHASHIKANT, C.V. SINGH, R.S. BARWAL and MANITA DANGI	151-157
Principal component analysis in production and reproduction traits of Frieswal cattle under field progeny testing OLYMPICA SARMA, R. S. BARWAL, C. V. SINGH, D. KUMAR, C. B. SINGH, A. K. GHOSH, B. N. SHAHI and S. K. SINGH	158-163
Degenerative renal pathology in swine: A comprehensive histopathological investigation in Rajasthan, India SHOBHA BURDAK, INDU VYAS, HEMANT DADHICH, MANISHA MATHUR, SHESH ASOPA, RENU	164-169
Evaluation of histopathological changes on acute exposure of profenofos in Swiss albino mice SONU DEVI, VINOD KUMAR, PREETI BAGRI and DEEPIKA LATHER	170-177
Temporal and spatial performance of rapeseed and mustard oilseed in India: A study in the context of Technology Mission on Oilseeds¹ LEKHA KALRA and S. K. SRIVASTAVA	178-190
Comparative economics of maize cultivation in major and minor maize producing districts of Karnataka – a study across farm size groups GEETHA, R. S. and S. K. SRIVASTAVA	191-203
A study on Usefulness of Participatory Newsletter for Potato growers in Udham Singh Nagar district of Uttarakhand RAMESH NAUTIYAL and ARPITA SHARMA KANDPAL	204–209
Training Needs of Hortipreneurs in Value Addition and fruit crop production in Kumaon Hills of Uttarakhand KRITIKA PANT and ARPITA SHARMA KANDPAL	210-215
Post-training Knowledge Assessment of the rural women about Mushroom Cultivation under TSP project, funded by ICAR ARPITA SHARMA KANDPAL, S. K. MISHRA and OMVEER SINGH	216-220
UAV Technology: Applications, economical reliance and feasibility in Indian Agriculture A. AJAY and S. SAI MOHAN	221-229

A study on dietary intake among school-going adolescent girls of Udaipur, Rajasthan during COVID-19

JYOTI SINGH^{1*} and NIKITA WADHAWAN²

¹Department of Food Science and Nutrition, College of Home Science, G.B. Pant University of Agriculture and Technology, Pantnagar-263145(U.S. Nagar, Uttarakhand), ²Department of Dairy and Food Technology, M.P.U.A.T, Udaipur (Rajasthan)

*Corresponding author's email id: 46182_jyotisingh@gbpuat-tech.ac.in

ABSTRACT: Adolescent girls deal with several major nutritional issues that impair their growth and development and their ability to support themselves as adults. The COVID-19 pandemic may have brought out changes in the lifestyle behaviour of adolescent girls, such as their food habits. Given this effect, a cross-sectional study was carried out to assess the dietary intake of adolescent girls aged 13 to 18 years in Udaipur, Rajasthan. Percent adequacy of dietary intake was assessed using the 24-hour dietary recall method and the average daily dietary intake was compared to the RDA (2020) and RDI (2010). The results of the study found that almost all the girls consumed very less amount of milk and its products, pulses, and GLV's along with cereals. However, there was adequate consumption of fruits, other vegetables, sugar, fats, and oils among both age groups. Intake of roots and tubers was adequate for girls aged 13 to 18 years but not for 16 to 18 aged girls. The average per adequacy nutrient intake of age group 13-15 and 16-18 years for energy, iron, calcium, vitamin B₁, B₂, B₃, and zinc was lower compared to RDA. Intake of CHO was found to be higher, whereas the intake of vitamin C, protein, fat, and folate was found to be adequate in the diet of both age groups. The findings of the study revealed that there was inadequate dietary intake among adolescent girls during the COVID-19 period, therefore it is important to promote the proper type of nutrition, and health education to avoid the development of unhealthy eating habits and thereby prevent the subsequent risk of degenerative diseases such as obesity, diabetes, cardiovascular pathologies, etc.

Key words: Adolescent girls, COVID-19, dietary intake, nutrition

In recent years, the coronavirus (COVID-19) outbreak has been regarded as the greatest threat to global health. This situation triggered a public health emergency of international significance, prompting the World Health Organization (WHO) to designate COVID-19 a pandemic, on March 11, 2020 (WHO, 2020). Many countries, including India, had to issue stay-at-home advisories and implement lockdown restrictions and social distancing procedures because of the global COVID-19 pandemic and the necessity to control the spread and transmission of the coronavirus (Ghosh *et al.*, 2020 and Patel *et al.*, 2021).

There have been numerous instances in the past where the public was subjected to restrictions as a result of natural catastrophes (such as tornadoes, hurricanes, and earthquakes) or man-made calamities (such as war and similar circumstances) have had serious implications, such as lack of resources and lifestyle changes, which has resulted

in disturbances in overall health of the population (Giorgadze *et al.*, 2011, Trevisan *et al.*, 1986, Ghosal *et al.*, 2020).

Adolescence is an important life stage that occurs between childhood and adulthood and is defined by significant physical, mental, and social changes (Patton *et al.*, 2016). Adolescent lifestyle activities, such as food consumption and physical exercise, create patterns that impact behaviour throughout adulthood and can have long-term health consequences (Blakemore and Mill, 2014 and Mokdad, 2016). In the early phases of infancy and adolescence, adequate nutrition is considered a possible factor for health (Glabaska *et al.*, 2020). It is critical to develop appropriate eating habits throughout this stage, i.e., the transition from childhood to maturity, to affect present health state and propensity to diseases in adulthood, such as obesity, diabetes, cardiovascular pathologies, and so on. Adolescent girls constitute an important segment

of the population. Their health status influences their reproductive functioning, pregnancy outcomes, birth weight, pregnancy wastage etc (Baliga *et al.*, 2014). The epidemic's isolation affects diet, especially among teenagers, who are more prone to developing bad eating habits (Singh and Wadhawan *et al.*, 2021). Nutritional intervention in combination with medical therapy, is a powerful tool to prevent and treat disease and disease-related symptoms. Emerging data from studies on COVID-19 and PASC, as well as studies on related respiratory viral infections and post-infection symptoms, indicates possible roles for nutrition in managing symptoms and improving clinical outcomes. By influencing both innate and adaptive immunity, nutrition may have an indirect impact on the risk of contracting COVID-19 (Kompaniyets *et al.*, 2021). Several globally recognized organisations have released dietary recommendations related to the SARS-CoV-2 pandemic (Rust and Ekmekcioglu, 2023). Various health organizations such as the WHO or UNICEF, as well as national organizations recommend the consumption of vegetables, fruits and whole grains, as the main components of a healthy diet (Montenegro-Landivar *et al.*, 2021). The Indian AYUSH ministry also made recommendations for food and nutrition during the confinement periods, as there is a strong link between the quality of a population's food and its health (Muscogiuri *et al.*, 2020).

A study conducted by Roso *et al.*, (2020) among adolescents of five different countries, including Spain, Italy, Brazil, Colombia, and Chile, to assess dietary trends and results of the study concluded that COVID-19 confinement had an impact on their eating habits where consumption of legumes, vegetables, and fruit intakes was significantly increased and fast-food intake was dramatically reduced. In another study by Pietrobelli *et al.* (2020), the results of the study discovered a rise in fruit consumption among children and adolescents in Italy.

It is necessary to evaluate the intake of food during COVID-19 among adolescent girls. Therefore, the study was designed to study the pattern of consuming

different food groups among adolescent girls residing in Udaipur, Rajasthan, India. This could help public health officials shape their suggestions for future measures in terms of nutrition policies for teenagers.

MATERIALS AND METHODS

The study was undertaken in March 2021. The target population was one hundred and twenty urban adolescent girls aged 13 to 18 years. The subjects were taken from two schools in Udaipur district, Rajasthan, and were selected by purposive random sampling method. The data were collected from respondents through a personal interview while keeping in mind the COVID-19 guidelines. A well-structured interview schedule was prepared and pretested to collect data on the dietary intake of all subjects. The dietary intake of adolescent girls was analyzed with the help of the 24-hour dietary recall method. Each participant was asked about the type and quantity of food consumed by them on the previous day. The type and quantity of food items consumed were assessed six times a day, including early morning, breakfast, mid-morning, lunch, evening snacks, and dinner. To maximize the accuracy of the study, standardized cup sets were used while interviewing them for their dietary intake throughout the whole day. Further converted in terms of raw ingredients and quantified raw food was reported in grams.

The mean dietary intake of different foods consumed was then calculated and compared with recommended dietary intake (NIN, ICMR, 2010) and recommended dietary allowances (NIN, ICMR, RDA, 2020) & for adolescent girls aged 13-15 and 16-18 years.

The data collected were tabulated and analysed using Diet Cal. Software. A z-test was used to compare nutrient and food intake with standard values.

RESULTS AND DISCUSSION

Food intake of adolescent girls: Intake of root & tubers (95.94 %), fruits (97.5%), sugar & jaggery

(92.6%), fats & oils (96.97 %), other vegetables (53.24%) among adolescent girls aged 13 to 15 years were found to be adequate as depicted in Table 1. In contrast, the consumption of food groups such as cereals (72.84%) was found to be lower than RDI and intake of pulses (41.35%), milk & milk products (44.05%), GLVs (46%), was found to be low than recommended dietary intake during COVID-19 period.

For girls aged 16 to 18 years, intake of fruits (91.42%), sugar & jaggery (91.12%), and fats & oils (99.03 %), and other vegetables (53.79 %), were near to the recommended values. In contrast, the consumption of cereals (81.48 %) was lower than RDI whereas intake of pulses (42.13%), milk & milk products (40.20%), GLVs (47.5%), roots & tubers (41.70 %) was not adequate as recommended by ICMR, NIN (2010) as presented in figure 2.

Nutrient Intake of adolescent girls: The results of the study found that the average per cent adequacy nutrient intake of age group 13-15 and 16-18 years for energy (65.86%, 64.43%), iron (40.44%, 41.77 %), calcium (43.79%, 36.96%), vitamin B₁ (71.00%, 73.76%), B₂ (30.80%, 29.78%), B₃ (46.79%, 47.06%), and zinc (62.79%, 63.39%) was low when

compared to RDA. Intake of CHO (177.58 %, 185.16 %) was found to be higher, whereas the intake of vitamin C (86.25%, 80.25%), protein (98.98%, 98.98%), fat (80.54%, 77.52%), and folate (85.40%, 81.51%) was found to be adequate in the diet of both age groups as shown in figure 3 and 4. The relationship between nutrition and the immune system is well known, so much attention is being paid to its role in COVID-19 (Butler *et al.*, 2021). Consuming a good quality diet is essential for a healthy immune system, and deficits in a variety of nutrients raise a person's risk of developing serious clinical symptoms as well as their vulnerability to viral infection. The purpose of this study was to analyse the dietary patterns of adolescent girls during the pandemic. The results of the present study reported that the daily nutrient intake of girls aged 13 to 18 years was not up to the mark as per recommendations. Intake of food groups such as root and tubers, fruits, vegetables, sugar and jaggery fats and oils was adequate whereas cereals, pulses, milk, and milk products was lower than RDI. Previous studies reported lower consumption of fruits and vegetables as per the recommendations among adolescents (Salwa *et al.*, 2021). The primary food groups in the diets of the selected teenagers were roots and tubers, followed by pulses, cereal and

Table 1: Daily mean food intake and per adequacy of food intake of adolescent girls (n=120)

Food groups	Age groups(years)	RDI Value	Actual intake Mean \pm SD	Z value	% Intake
Cereals & millets	13-15 years	330	240.38 \pm 37.82	2.36*	72.84
	16-18 years	330	268.89 \pm 29.90	2.04*	81.48
Pulses	13-15 years	60	24.81 \pm 12.5	2.81**	41.35
	16-18 years	75	31.6 \pm 10.56	4.10**	42.13
Milk & milk products	13-15 years	500 ml	220.26 \pm 113.77	2.45*	44.05
	16-18 years	500 ml	201 \pm 90.98	3.28**	c
Roots & tubers	13-15 years	100	95.94 \pm 48.69	0.08 ^{NS}	95.94
	16-18 years	200	83.41 \pm 33.13	3.51**	41.70
Green leafy vegetables	13-15 years	100	46 \pm 20.28	2.66**	46
	16-18 years	100	47.5 \pm 9.57	5.48**	47.5
Other vegetables	13-15 years	200	106.49 \pm 65.28	1.43 ^{NS}	53.24
	16-18 years	200	107.56 \pm 47.51	1.94 ^{NS}	53.79
Fruits	13-15 years	100	97.5 \pm 24.92	0.10 ^{NS}	97.5
	16-18 years	100	91.42 \pm 22.67	0.37 ^{NS}	91.42
Sugar and jaggery	13-15 years	25	23.15 \pm 1.23	1.50 ^{NS}	92.6
	16-18 years	25	22.78 \pm 1.09	1.24 ^{NS}	91.12
Fat/oil (visible)	13-15 years	40	38.79 \pm 2.78	0.43 ^{NS}	96.97
	16-18 years	35	34.66 \pm 5.32	0.003 ^{NS}	99.031

*Significant at (p<0.05), **Significant at (p<0.01), NS= non-significant, # Source: RDI- Recommended dietary intake, NIN, ICMR (2010)

Table 2: Mean nutrient intake and per adequacy of nutrient intake of the subjects (n=120)

Nutrients	Age groups	Reference Value #	Actual Intake Mean SD	Z value	% Intake
Energy(Kcal/d)	13-15 years	2400	1580.80±237.46	3.4**	65.86
	16-18 years	2500	1610.79±230.07	3.86**	64.43
Protein(g/d)	13-15 years	43.2	42.76±7.22	0.06 ^{NS}	98.98
	16-18 years	46.2	45.75±7.70	0.05 ^{NS}	99.03
Total fat(g/d)	13-15 years	66.66	53.69±7.99	1.62 ^{NS}	80.54
	16-18 years	66.66	51.68±8.77	1.70 ^{NS}	77.52
CHO(g/d)	13-15 years	130	230.86±25.70	-3.92**	177.58
	16-18 years	130	240.72±28.56	-3.87**	185.16
Ca(mg/d)	13-15 years	1000	437.96±176.94	3.17**	43.79
	16-18 years	1050	388.13±181.51	3.64**	36.96
Fe(mg/d)	13-15 years	30	12.13±2.06	8.67**	40.44
	16-18 years	32	13.26±1.75	10.65**	41.77
Vit C(mg/d)	13-15 years	66	56.92±26.03	0.34 ^{NS}	86.25
	16-18 years	68	54.57±24.97	0.53 ^{NS}	80.25
B1(mg/d)	13-15 years	1.6	1.13±0.16	2.93**	71.00
	16-18 years	1.7	1.25±0.14	3.21**	73.76
B2(mg/d)	13-15 years	2.2	0.67±0.18	8.5**	30.80
	16-18 years	2.3	0.68±0.15	10.51**	29.78
B3(mg/d)	13-15 years	16	7.48±1.2	7.1**	46.79
	16-18 years	17	8±1.12	8.03**	47.06
Folate(µg/d)	13-15 years	245	209.25±54.72	0.65 ^{NS}	85.40
	16-18 years	270	220.09±57.54	0.86 ^{NS}	81.51
Zn(mg/d)	13-15 years	12.8	8.03±1.17	4.07**	62.79
	16-18 years	14.2	9±1.25	4.13**	63.39

*Significant at (p<0.05), **Significant at (p<0.01), NS=non-significant, # Source: RDA- Recommended dietary Allowance, NIN, ICMR (2020)

millet, and sugar; there was a modest amount of fats and oils and vegetables, but a dearth of green leafy vegetables, milk and milk products, fruits, meat, and poultry when compared to the recommended dietary allowances (Seth, 2021). Bharathi, 2021 concluded that rural women of all ages did not consume at least a third of the RDI for pulses. Compared to rural women, urban women have improved intake but still consume 13 to 37% less.

Nutrient intake among adolescent girls in the present study was found to be inadequate. Intake of energy, iron, calcium, vitamin B1, B2, B3, and zinc was low when compared to RDA. Consumption of carbohydrates was found to be higher, whereas the intake of vitamin C, protein, fat and folate was adequate in their diet. Previous studies reported a deficient supply of energy, protein, and micronutrients such as iron, calcium, vitamin B12, folate, calcium, and zinc among Indian adolescent girls (Maliye *et al.*, 2010, Sachan *et al.*, 2013, Doustmohammadian *et al.*, 2013, Gupta *et al.*, 2018, Solanki *et al.*, 2021, Sari *et al.*, 2022).

During the COVID-19 pandemic, Vyas and Rawal, 2022 reported that the consumption of pulses, fruits, and vegetables among adolescents of the age group 12 to 14 years was found to increase and they also reported that most (81%) adolescents included milk in their diet daily which is contrary to the present study. In a representative German online survey, children, and adolescents (1–14 years old) showed an increase in healthier behaviours, such as increased fruit and vegetable consumption (Koletzko *et al.*, 2021). In a study conducted on Canadian university students, it was reported that the nutrient and caloric intake were significantly reduced during the pandemic, alcohol intake increased significantly whereas the frequency of consuming food groups across the board decreased during the pandemic (Butler and Barrientos, 2021). 10% of Belgian participants often could not afford to eat a healthy diet during the pandemic period (Pfeifer *et al.*, 2021). Fruit and vegetable consumption decreased and the consumption of soft drinks, sweets, bread and salty snacks increased (Vandevijvere, *et al.*, 2020, Huber




Fig. 1: Percent adequacy of food groups among adolescent girls (13-15 years)




Fig. 2: Percent adequacy of food groups among adolescent girls (16-18 years)




Fig. 3: Percent adequacy of nutrients among adolescent girls (13-15 years)




Fig. 4: Percent adequacy of nutrients among adolescent girls (16-18 years)

et al., 2020, Zarah *et al.*, 2020). Adolescents reported high consumption of food rich in energy as rice, pasta, cheese, or legumes, a lower consumption of protein than three servings per week and deficient intake of vitamins and minerals (Martín *et al.*, 2022). Previous studies among adolescents on dietary intake showed dietary inadequacy. Adolescent's dietary consumption and food choices may be influenced by a variety of factors, including household economic status, uneven intra-familial food distribution, food taboos, dietary restrictions during menstruation, peer pressure, food choices, culture, mass media, and body image perception (Melaku *et al.*, 2018, Dalky *et al.*, 2016, Ethiopian Federal Ministry of Health, 2016). COVID-19 resulted in both favorable and unfavorable changes in dietary pattern among adolescent. The present study showed better dietary intake than pre-COVID time, but during COVID, the consumption was also not up to the mark as recommended by RDI and RDA. Reasons for reduced dietary intake predominately

included limited availability and increased price, and there were associations with poor food choices and mental health conditions including depression and anxiety, and sedentary time (Sidor and Rzymiski, 2020).

CONCLUSION

During adolescence, adequate nutrition is a potential health factor. COVID-19 has an impact on one's lifestyle, particularly nutrition. The results of this study provided the first description of the dietary intake of adolescent girls of age groups 13 to 18 years during the outbreak period. The key findings of the study stated that dietary intake was inadequate among girls during COVID-19 which indicated inappropriate dietary consumption among adolescent girls. The impact of social isolation and the pandemic on eating behaviour during COVID-19 should not be underestimated as it has already had acute effects and will likely produce long-term deleterious effects

on health as well. It is a matter of the utmost importance to abandon the negative habits acquired by the adolescents, therefore, families of the subjects, as well as nutritional and health professionals, are urged to educate and encourage these adolescents to improve their dietary habits. Understanding the adolescent's nutritional behaviour through our study will help public health authorities to modify future regulations on adolescents' nutritional recommendations.

REFERENCES

- Baliga, S., S., Naik, V., A. and Mallapur, M., D. (2014). Nutritional status of adolescent girls residing in rural area: A community-based cross-sectional study. *Journal of the Scientific Society*, 41(1): 22-2.
- Bertrand, L., Shaw, K., A., Ko, J., Deprez, D., Chilibeck, P.D. and Zello, G.A. (2021). The impact of the coronavirus disease 2019 (COVID-19) pandemic on university students' dietary intake, physical activity, and sedentary behaviour. *Applied Physiology, Nutrition, and Metabolism*, 15: 1-8.
- Bharathi, R. (2021). Dietary and Nutrient Intakes of Rural and Urban Women: A Study from South India. 2021. *International Journal of Theoretical and Applied Sciences*, 13(1): 19-25.
- Blakemore, S. J. and Mills, K. L. (2014). Is adolescence a sensitive period for sociocultural processing? *Annual Review of Psychology*, 65: 187-207.
- Butler, M.J. and Barrientos, R.M. (2021). The impact of nutrition on COVID-19 susceptibility and long-term consequences. *Brain Behaviour and Immunity*, 87:53-54.
- Dalky, H. F. and Al-drabaah, T. K. (2016). Eating habits and associated factors among adolescent students in Jordan. *Clinical nursing research*, 26: 538-52.
- Doustmohammadian, A. Keshavarz, S., A., Doustmohammadian, S., Abtahi, M. and Shahani, M. (2013). Nutritional status and dietary intake among adolescent girls. *Journal of Paramedical Sciences (JPS)*, 4: 72-77.
- Ethiopian Federal Ministry of Health (2016). National Guideline on adolescent. Addis Ababa: Maternal Infant and Young Child Nutrition.
- Ghosal, S., Sinha, B., Majumder, M. and Misra, A. (2020). Estimation of effects of nationwide lockdown for containing coronavirus infection on worsening of glycosylated haemoglobin and increase in diabetes-related complications: A simulation model using multivariate regression analysis. *Diabetes Metabolic Syndrome*, 14: 319-23.
- Ghosh, A., Nundy, S. and Mallick, T. K. (2020). How India is dealing with COVID-19 pandemic. *Sensors*, 1:100021.
- Giorgadze, T., Maisuradze, I., Japaridze, A., Utiashvili, Z. and Abesadze G. (2011). Disasters and their consequences for Public Health. *Georgian Medical News*, 194: 59-63.
- Glabska, D., Guzek, D., Groele, B. and Gutkowska, K. (2020). Fruit and vegetables intake in adolescents and mental health: A systematic review. *Rocz. PaDstwowej ZakBadu Hig.*, 71: 15-25.
- Gupta, A., Noronhab, J. A., Shobhab and Garg, M. (2018). Dietary intake of macronutrients and micronutrients among adolescent girls: A cross-sectional study. *Clinical Epidemiology and Global Health*, 6: 192-197.
- Huber, B.C., Steffen, J., Schichtiger, J. and Brunner, S. (2020). Altered nutrition behaviour during COVID-19 pandemic lockdown in young adults. *European Journal of Nutrition*, 1: 1-10.
- Koletzko, B., Holzapfel, C., Schneider, U. and Hauner, H. (2021). Lifestyle and Body Weight Consequences of the COVID-19 Pandemic in Children: Increasing Disparity. *Annals of Nutrition and Metabolism*, 77: 1-3.
- Kompaniyets L., Goodman A.B., Belay B., Freedman D.S., Sucusky M.S., Lange S.J., Gundlapalli A.V., Boehmer T.K. and Blanck H.M. (2021). Body Mass Index and Risk for

- COVID-19–Related Hospitalization, Intensive Care Unit Admission, Invasive Mechanical Ventilation, and Death—United States. *Morbidity and Mortality Weekly Report*, 70: 355–361.
- Maliye, C.H, Deshmukh, P., Gupta, S., Kaur, S., Mehendale, A. and Garg, B. (2010). Nutrient intake amongst rural adolescent girls of Wardha. *Indian Journal of Community Medicine: official publication of Indian Association of Preventive & Social Medicine*, 35(3): 400–402.
- Martín-R., A., Tornero-Aguilera, J. F., López-Pérez, P. J. and Clemente-Suárez, V. J. (2022). Dietary patterns of adolescent students during the COVID-19 pandemic lockdown. *Physiology & behavior*, 249: 113764.
- Melaku, Y., Dirar, A., Feyissa, G. T. and Tamiru, D. (2018). Optimal dietary practices and nutritional knowledge of school adolescent girls in Jimma town, south West Ethiopia. *International Journal of Adolescence and Youth*, 23:299–307.
- Mokdad, A. H., Forouzanfar, M. H., Daoud, F., Mokdad, A. A., El Beheraoui, C., Moradi-Lakeh, M., Kyu, H. H., Barber, R. M., Wagner, J. and Cercy, K. (2016). Global burden of diseases, injuries, and risk factors for young people's health during 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*, 387: 2383–2401.
- Montenegro-Landivar, M.F., Tapia-Quiros, P., Vecino, X., Reig, M., Valderrama, C., Granados, M., Cortina, J.L. and Saurina, J. (2021). Polyphenols and their potential role to fight viral diseases: An overview. *Science of the Total Environment*, 801: 149719.
- Muscogiuri, G., Barrea, L., Savastano, S. and Colao, A. (2020). Nutritional recommendations for COVID-19 quarantine. *European Journal of Clinical Nutrition*, 74: 850–851.
- NIN. ICMR. (2010). Dietary Guidelines for Indians: A Manual of National Institute of Nutrition, Hyderabad.
- Patel, A., Agarwal, R., Rudramurthy, S. M., Shevkani, M., Xess, I., Sharma, R., Savio, J., Sethuraman, N., Madan, S., Shastri, P., Thangaraju, D., Marak, R., Tadepalli, K., Savaj, P., Sunavala, A., Gupta, N., Singhal, T., Muthu, V., Chakrabarti, A. and MucoCovi. (2021). Multicenter Epidemiologic Study of Coronavirus Disease-Associated Mucormycosis, India. *Emerging Infectious Diseases*, 27(9): 2349–2359.
- Patton, G. C., Sawyer S. M., Santelli, J. S., Ross, D. A., Afifi, R., Allen, N. B., Arora, M., Azzopardi, P., Baldwin, W. and Bonell, C. (2016). Our future: a Lancet commission on adolescent health and wellbeing. *Lancet*, 387: 2423–2478.
- Pfeifer, D., Rešetar, J., Kljusuri, J.G., Krbavi, I.P., Bender, D.V., Rodríguez-Pérez, C., Ruíz-López, M.D. and 'atali, Z. (2021). Cooking at home and adherence to the Mediterranean diet during COVID-19 confinement: The experience from the Croatian COVIDiet Study. *Frontier Nutrition*, 8: 617721.
- Pietrobelli, A., Pecoraro, L., Ferruzzi, A., Heo, M., Faith, M., Zoller, T., Antoniazzi, F., Piacentini, G., Fearnbach, S. N. and Heymsfield, S. B. (2020). Effects of COVID-19 Lockdown on Lifestyle Behaviors in Children with Obesity Living in Verona, Italy: A Longitudinal Study. *Obesity*, 28(8): 1382–1385.
- Roso, R., M.B., de Carvalho Padilha P., Mantilla-Escalante, D. C., Ulloa, N., Brun, P., Acevedo-Correa, D., Arantes Ferreira Peres, W., Martorell, M., Aires, M. T., de Oliveira Cardoso, L., Carrasco-Marín, F., Paternina-Sierra, K., Rodriguez-Meza, J. E., Montero, P. M., Bernabè, G., Pauletto, A., Taci, X., Visioli, F. and Dávalos, A. (2020). Covid-19 Confinement and Changes of Adolescent's Dietary Trends in Italy, Spain, Chile, Colombia and Brazil. *Nutrients*, 12(6):1807.
- Rust, P. and Ekmekcioglu, C. (2023). The Role of Diet and Specific Nutrients during the COVID-19 Pandemic: What Have We Learned over the Last Three Years? *International Journal of Environmental*

- Research and Public Health*; 20(7):5400.
- Sachan, B., Idris, M.Z., Jain, S., Kumari, R. and Singh, A. (2013). Nutrient Intake Among School-Going Adolescent Girls in Lucknow district, India. *Journal of Biomedical and Pharmaceutical Research*, 2(4): 101-105.
- Salwa, M., Subaita, F., Choudhury, S. R., Khalequzzaman, M., Al Mamun, M. A., Bhuiyan, M. R. and Haque, M. A. (2021). Fruit and vegetables consumption among school-going adolescents: Findings from the baseline survey of an intervention program in a semi-urban area of Dhaka, Bangladesh. *PloS one*, 16(6): e0252297.
- Sari, P., Herawati, D. M. D., Dhamayanti, M. and Hilmanto, D. (2022). The Study of Nutrient Intake and Adolescent Girls' Quality of Life in a Rural Area of Indonesia. *Children (Basel, Switzerland)*, 9(8): 1248.
- Seth U. (2021). Eating Behavior and Nutritional Status of Adolescents in Peri-Urban Delhi. *Indian Journal of Nutrition*, 8(1): 225.
- Sidor, A. and Rzymiski, P. (2020). Dietary choices and habits during COVID-19 lockdown: experience from Poland. *Nutrients*, 12:1657.
- Singh, J. and Wadhawan, N. (2021). Effect of Covid-19 on Food Consumption Pattern of Indian Adolescent Girls. *Acta Scientific Nutritional Health*, 5(10): 77-87.
- Solanki, S., and Solanki, M. (2021). A study to determine calorie and protein intake in adolescent school girls of Indore city, M.P., India. *International Journal of Medical Research and Review*, 9(6): 406-411.
- Trevisan, M., Celentano, E., Meucci, C., Farinaro, E., Jossa, F. and Krogh, V. (1986). Short-term effect of natural disasters on coronary heart disease risk factors. *Arteriosclerosis*, 6: 491-4.
- Vandevijvere, S., De Ridder, S., Drieskens, S., Charafeddine, R., Berete, F. and Demarest, S. (2020). Food insecurity and its association with changes in nutritional habits among adults during the COVID-19 confinement measures in Belgium. *Public Health Nutrition*, 9: 1-12.
- Vyas, M. and Rawal, S. (2022). Covid 19 pandemic and its effect on the Nutritional status of adolescents- a study in Surat. *IJARIE*, 8(10): 334-342.
- World Health Organization (WHO) Geneva: 2020. Accessed on 20 Nov 2023. WHO Director-General's Opening Remarks at the Media Briefing on COVID-19. Accessed from: <https://www.who.int/director-general/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19—11-march-2020>.
- Zarah, A.B., Enriquez-Marulanda, J. and Andrade, J.M. (2020). Relationship between dietary habits, food attitudes and food security status among adults living within the United States three months post-mandated quarantine: A cross-sectional study. *Nutrients*, 12: 3468.

Received: January 23, 2024

Accepted: April 07, 2024