

Print ISSN : 0972-8813  
e-ISSN : 2582-2780

[Vol. 24(1) January-April 2026]

# Pantnagar Journal of Research

(Formerly International Journal of Basic and  
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology  
Pantnagar, U.S. Nagar; Uttarakhand, Website : [gbpuat.res.in/PJR](http://gbpuat.res.in/PJR)



## ADVISORY BOARD

### Patron

**Prof. Shivendra Kumar Kashyap**, Ph.D., Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

### Members

**Prof. S. K. Verma**, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. Jitendra Kwatra**, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. S.S. Gupta**, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. D. Kumar**, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. Alka Goel**, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. R.S. Jadoun**, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. Lokesh Varshney**, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. Avdhesh Kumar**, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. Subhash Chandra**, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Prof. Ramesh Chandra Srivastava**, Ph.D., Dean, College of Basic Sciences & Humanities, G.B.P.U.A.T., Pantnagar, India

## EDITORIAL BOARD

### Members

**A.K. Misra**, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

**Anand Shukla**, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

**Anil Kumar**, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

**Ashok K. Mishra**, Ph.D., Kemper and Ethel Marley Foundation Chair, W.P. Carey Business School, Arizona State University, U.S.A.

**Binod Kumar Kanaujia**, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

**D. Ratna Kumari**, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

**Deepak Pant**, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

**Desirazu N. Rao**, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

**G.K. Garg**, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Humnath Bhandari**, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

**Indu S. Sawant**, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

**Kuldeep Singh**, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

**Muneshwar Singh**, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

**Omkar**, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

**P.C. Srivastav**, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

**Prashant Srivastava**, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

**Puneet Srivastava**, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

**R.K. Singh**, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

**Ramesh Kanwar**, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

**S.N. Maurya**, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agri. & Tech., Pantnagar, India

**Sham S. Goyal**, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

**Umesh Varshney**, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

**V.D. Sharma**, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

**V.K. Singh**, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

**Vijay P. Singh**, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

**Nikitin Georgy**, Ph.D., Vice-Rector cum Professor St. Petersburg State University of Veterinary Medicine St. Petersburg, Russia

**Sanjay Jambhulkar**, Ph.D., Ex-Scientific Officer 'H' Head Mutation Breeding Section Bhabha Atomic Research Centre, Mumbai Professor, Homi Bhabha National Institute, Mumbai

**Vijay Kubsad**, Ph.D., Adjunct Professor Washington State University USA

**Kiran Tota-Maharaj**, Ph.D., Professor Royal Agricultural University (RAU) Cirencester, Gloucestershire GL76JS England, UK

**Ravindra P. Veeranna**, Ph.D., Dean, Xavier University, College of Veterinary Sciences, Aruba

**Javed Akhtar**, Ph.D., University of Technology and Applied Sciences, Muscat

**Basant Maheshwari**, Ph.D., Western Sydney University, Australia

**Vijay P. Singh**, Ph.D. Professor Emeritus, Texas A&M University College station, Texas, USA

**Sanjay Sharma**, Ph.D. Director, Uttarakhand Council of Biotechnology

**Krishna Nemali**, Ph.D. Controlled Environment Agriculture Extension Specialist

### Editor-in-Chief

**K.P. Raverkar**, Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

### Managing Editor

**S.K. Guru**, Ph.D., Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

### Assistant Managing Editor

**Jyotsna Yadav**, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

### Technical Manager

**S.D. Samantaray**, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

### Development

**Dr. S.D. Samantaray**, Professor & Head

**Brijesh Dumka**, Developer & Programmer

## CONTENTS

<b>Indian mustard: Phytoremediation tool for soil health management (Review)</b>	<b>1</b>
M.L. DOTANIYA, C.K. DOTANIYA, KULDEEP KUMAR, JAKIR HUSSAIN, R.K. DOUTANIYA, H.M. MEENA and MANJU LATA	
<b>From profile to problems: A study on socio-economic status and constraints of rural women in dairy farming</b>	<b>14</b>
ADITI PATHAK and ARPITA SHARMA KANDPAL	
<b>Diversification and intensification in rice–wheat cropping system for enhancing productivity under Tarai zone of Uttarakhand</b>	<b>20</b>
PESALA SANTHA KUMAR, DINESH KUMAR SINGH, SUBHASH CHANDRA, ROHITASH SINGH and AJEET PRATAP SINGH	
<b>Sensor based nitrogen application impact on productivity, economics and nutrient uptake in maize</b>	<b>25</b>
AMIT BHATNAGAR and AJAY KUMAR	
<b>Association of germination and seedling vigour attributes with green forage yield in oat (<i>Avena sativa</i> L.) genotypes through correlation and path coefficient analysis</b>	<b>31</b>
DEVYANI RAMOLA, AKRITI BALLABH SANSKRITI JOSHI, MANISH MEHRA and BIRENDRA PRASAD	
<b>Genetic divergence and yield trait impact in French bean (<i>Phaseolus vulgaris</i> L.)</b>	<b>43</b>
BHAWANA MAMGAIN, B.P. NAUTIYAL, S.C. PANT, VAISHALI THAPA, HEIGRUJAM RIYASANA and MAYANK	
<b>Arabic-gum based edible coatings preserving the post-harvest quality of Guava fruits</b>	<b>48</b>
ESHA TEWARI and OMVEER SINGH	
<b>Efficacy of catechol and plant growth regulators on the vegetative growth and flowering of African Marigold (<i>Tagetes erecta</i> L. cv. Pusa Narangi Gainda)</b>	<b>59</b>
SURYANSH SRIVASTAVA, AFSAR AHMAD, MD. RAMJAN and KHAN BILAL MUKHTAR AHMED	
<b>Host response of diverse pulse genotypes to root-knot nematode (<i>Meloidogyne incognita</i>)</b>	<b>67</b>
SAROJ YADAV, EKTA RATHORE and ANIL KUMAR	
<b>Effect of egg size on egg quality traits, fertility, hatchability and day-old chick weight</b>	<b>75</b>
ISLAM UDDIN SHEIKH, AZMAT ALAM KHAN, ARNAB JYOTI KALITA, SHEIKH ADIL HAMID, IRFAN AKRAM BABA, HENNA HAMADANI, BUSHRA ZAFFER and INSHA AFZAL	
<b>Pathomorphological studies on verminous pneumonia in a Gaddi goat</b>	<b>81</b>
SWASTI SHARMA, SONALI MISHRA and R.D. PATIL	
<b>Pathological findings in congestive heart failure in dog: A case report</b>	<b>85</b>
MUNISH BATRA and STUTI SINGH	

<b>Histopathological and haematological evaluation of the ameliorative effect of piperine on quinalphos-induced subacute toxicity in Swiss Albino Mice</b> PREETI SINGH, VINOD KUMAR, PREETI BAGRI and DEEPIKA LATHER	89
<b>Phytochemical screening and antioxidant activities of locally available medicinal plants</b> PREETI BAGRI, VIJEYTA TIWARI, ARCHANA LOHIYA and VINOD KUMAR	99
<b>Natural Farming Inputs (NFIs) as sustainable alternatives for enhancing the growth and development of Chickpea (<i>Cicer arietinum</i> L.)</b> SHAIFALI and A.K. SHARMA	108
<b>Integrating Bayesian Calibration, Hierarchical Spatial Regression, and Structural–Temporal Models to Explain and Predict Wheat Yield in the Tarai–Bhabar Belt of Uttarakhand, India</b> NEHA BHATT and VINOD KUMAR	114
<b>Phytochemical characterization and immuno-antioxidant potential of <i>Moringa oleifera</i> leaf extract in chicken lymphocytes</b> ANAMITRA RUJ, SONU AMBWANI and LATA PALIWAL	128
<b>Biochemical changes associated with triclosan exposure in the freshwater fish <i>Cyprinus carpio</i></b> MADHU SHARMA, YASHPAL, RIJUL RANA, DEEKSHA, DEEPIKA THAKUR and TARANG KUMAR SHAH	140
<b>Assessment of awareness among respondents towards Pradhan Mantri Jan Dhan Yojana: A study in Haryana</b> MEGHA GOYAL, SUMAN GHALAWAT, MINAKSHI, ATUL DHINGRA and MINI	148
<b>Youth perception and constraints in agri-entrepreneurship: A factor analytic study</b> JYOTI, SUMAN GHALAWAT, MEGHA GOYAL, NEELAM KAUSHAL, SUNIL KUMAR and VIKRANT HOODA	156
<b>Design of Object Detection Model Using Deep Learning for Autonomous Driving in Adverse Weather Conditions: A GAN-Based Restoration and Depth-Aware Fusion Framework</b> MUMUKSHU BHATT, SUBODH PRASAD, RAJESH SINGH, ASHOK KUMAR and POOJA TAMTA	165
<b>A novel mechanical purification approach for <i>Grewia optiva</i> Fibre-Reinforced epoxy composites: Effect of fibre geometry on structural performance</b> DEEPA SINGH, NEERAJ BISHT, TARUN KUMAR SAXENA and KAMAL RAWAT	172
<b>Evaluating Long-Term rainfall and temperature trends across the Shipra River basin</b> POONAM SINGH, MOHAN LAL, DHEERAJ KUMAR, P. K. SINGH and GOVIND VERMA	184
<b>Rainfall-runoff modelling using soft computing techniques for various watersheds of Madhya Pradesh, India</b> SEEMA KUSHWAHA and PRAVENDRA KUMAR	196
<b>Prevalence and associated risk factors of functional constipation among the selected adult population of Pantnagar, U.S. Nagar, Uttarakhand, India</b> JYOTI SINGH, NEETU DOBHAL, ARCHANA KUSHWAHA, N.C SHAHI and SABBU SANGEETA	204
<b>Tracking nutritional transitions: A comparative study of child malnutrition (0–5 years) trends across districts of Madhya Pradesh</b> SHALINI CHAKRABORTY	212
<b>Determinants influencing tourist selection of rural tourism destinations using Garrett Ranking Technique: Evidence from Supi village, Uttarakhand</b> NEHA PANDEY and KAMAL UPRETI	221

## Indian mustard: Phytoremediation tool for soil health management

M.L. DOTANIYA<sup>1\*</sup>, C.K. DOTANIYA<sup>2</sup>, KULDEEP KUMAR<sup>3</sup>, JAKIR HUSSAIN<sup>4</sup>, R.K. DOUTANIYA<sup>5</sup>, H.M. MEENA<sup>6</sup> and MANJU LATA<sup>7</sup>

<sup>1</sup>ICAR-Indian Institute of Rapeseed-Mustard Research, Bharatpur-321303, <sup>2</sup>Department of Soil Science & Agricultural Chemistry, SKRAU, Bikaner-334006, <sup>3</sup>ICAR-Indian Institute of Soil & Water Conservation, Dehradun, Research Centre, Kota-324 002, <sup>4</sup>Central Water Commission, Department of Water Resources, River Development & Ganga Rejuvenation (DoWR, RD & GR), Ministry of Jal Shakti, New Delhi, <sup>5</sup>Department of Agronomy, SKN College of Agriculture, Jobner- 303328, <sup>6</sup>ICAR - Central Arid Zone Research Institute, Jodhpur-342003, <sup>7</sup>University of Rajasthan, Jaipur-302004

\*Corresponding author's email id: mohan30682@gmail.com

**ABSTRACT:** Metal and metallic dynamics in soil affect the ecosystem services and mediated the food chain contamination over a period. Industrialization of a country boost the economic growth, whereas waste generation during the different process poorly manage will be the source of pollution. Dumping of industrial effluents in natural resources reach the human body via food chain contamination is a major challenges in-front of researcher and policy makers. Different type of remediation methods are practicing, among them, phytoremediation is an eco-friendly and cheaper. Selection of Indian mustard (*Brassica juncea* L.) for phytoremediation is well established and popular practice in many countries. It is having the higher biomass and poor translocation of heavy metals from shoot to edible part. Research and development activities have been improved the greater biomass and crop yield in metal contaminated soils also. Long- term cultivation of mustard crop on contaminated soil improved the soil health parameters and reduces the metal concentration in soil solution. Use of mustard crop as a phytoremediation is a good practice without compromising the agricultural crop production in metal toxicity areas.

**Keywords:** Food chain contamination, Heavy metal, Indian mustard, Remediation methods, Soil health

Metal and metallic dynamics in soil greatly influenced by the origin of metals and their dispersion in ecosystem (Donald, 2003). Advancement of scientific tool and techniques explored the extraction and purification of metals from ore and further utilization of metal for human welfare. However, human development also generating the waste, mighty recovers the metal and discharge in natural resources (Gupta *et al.*, 2012). Soil contamination poses a significant threat to global agriculture, ecosystems, and human health (Dehkordi *et al.*, 2024). Heavy metals such as zinc (Zn), cadmium (Cd), arsenic (As), lead (Pb), mercury (Hg), chromium (Cr) and copper (Cu) accumulate in soils due to industrial activities, mining, wastewater irrigation, and excessive fertilizer use (Singh *et al.*, 2011; Dotaniya *et al.*, 2016a). These contaminants reduce soil fertility, inhibit plant growth, and enter the food chain, leading to bioaccumulation and toxicity (Dotaniya *et al.*, 2022b). Traditional remediation methods, like soil excavation or chemical treatments, are often

expensive, disruptive, and environmentally harmful. Phytoremediation emerges as a sustainable, cost-effective alternative, utilizing plants to extract, stabilize, or degrade pollutants in soil (Rajendiran *et al.*, 2018).

Phytoremediation involves several mechanisms: phytoextraction (uptake and accumulation of contaminants in harvestable plant parts), phytostabilization (immobilization in roots to prevent leaching), rhizofiltration (filtration by roots in aqueous environments), and phytovolatilization (conversion to volatile forms) (Sharma and Dubey, 2005). Among hyperaccumulator plants—species capable of accumulating metals at levels 100 times higher than average—Indian mustard (*Brassica juncea*) stands out for its rapid growth, high biomass production, and tolerance to heavy metals (Ojuederie *et al.*, 2022). Native to India and widely cultivated for oilseed, this plant has been extensively studied for its ability to remediate metal-contaminated soils, thereby improving soil health by reducing toxicity and restoring nutrient balance. Indian mustard's role

in phytoremediation is particularly relevant in regions like India and China, where agricultural lands are increasingly polluted. Studies highlight its efficiency in accumulating metals like Zn and Cd, making it a promising tool for eco-friendly soil restoration (Shiyab *et al.*, 2009). This draft explores the metal toxicity & its mechanism in soil-plant continuum, phytoremediation potential of Indian mustard, phytoremediation mechanisms, enhancement techniques, benefits to soil health and challenges.

### ***Role of metal in plant nutrition***

As per the criteria of essentiality described by the Arnon and Stout (1939) indicated that many metal are playing role in the plant nutrition in required quantity however, exceed the concentration in soil or plant showed as toxicity system. Metallic micronutrients are essential for higher plants, functioning primarily as cofactors in enzymes, structural components in proteins, and participants in redox reactions (Kaur and Garg, 2021). They enable biochemical processes that macronutrients alone cannot support, particularly those involving electron transfer, oxygen handling, and specialized catalysis. Unlike zinc ( $Zn^{2+}$ ), which does not change oxidation state and acts mainly as a Lewis acid (electrophile activator), the others (Fe, Mn, Cu, Mo, Ni) are redox-active transition metals. This allows them to shuttle electrons, cycle between oxidation states ( $Fe^{2+} \rightleftharpoons Fe^{3+}$ ), and catalyze reactions with high specificity due to unique coordination geometries, ligand preferences, and redox potentials. Table 1 listed the role of essential metal in plant nutrition.

### ***Heavy metal toxicity***

Metal toxicity in the soil ecosystem refers to the harmful effects caused by elevated concentrations of metals (especially heavy metals and metalloids) that exceed natural background levels or plant/organism tolerance thresholds (Minhas *et al.*, 2021). While, some metals (Fe, Mn, Zn, Cu, Mo, Ni) are essential micronutrients at low levels, they become toxic at high concentrations. Non-essential metals (Cd, Pb, As, Hg, Cr) are toxic even at trace amounts (Table 2). These metals persist in soils due to their non-biodegradable nature, accumulate over time, and affect the entire soil food web: plants, soil microbes, invertebrates, and higher trophic levels, ultimately threatening agriculture, biodiversity, food safety, and human health *via* the food chain.

### ***Effect of metal toxicity in the soil ecosystem***

Toxic substances in soil ecosystems affect organisms and ecological processes through several biological and chemical mechanisms. Contaminants such as heavy metals, pesticides, industrial chemicals, and persistent organic pollutants can cause direct toxicity to soil microorganisms by damaging cell membranes, proteins, and DNA, thereby reducing microbial growth and activity (Sparks, 2003). Many pollutants also cause enzyme inhibition, where toxic metals like cadmium, mercury, and lead bind to enzyme active sites or sulfhydryl groups, disrupting essential biochemical reactions involved in nutrient cycling. Another important mechanism is oxidative stress, in which toxic compounds stimulate the formation of reactive oxygen species (ROS) that damage

**Table 1: Essential plant nutrients and their functions**

Metal	Uptake form	Function
Iron (Fe)	$Fe^{2+}$ (ferrous), though $Fe^{3+}$	Chlorophyll biosynthesis, Photosynthesis, Nitrogen metabolism, electron flow and preventing oxidative damage
Manganese (Mn)	$Mn^{2+}$	Photosynthesis and enzyme activation, Photosystem II water-splitting complex
Zinc (Zn)	$Zn^{2+}$	Structural and catalytic in >200–300 enzymes/proteins
Copper (Cu)	$Cu^{2+}$	Redox catalysis in oxidative enzymes
Molybdenum (Mo)	$MoO_4^{2-}$	Nitrogen assimilation/fixation
Nickel (Ni)	$Ni^{2+}$	Urea metabolism

cellular components such as lipids, proteins, and nucleic acids in soil organisms (Pepper *et al.*, 2014). Toxic chemicals can also undergo bioaccumulation, where substances accumulate in organisms such as earthworms, nematodes, and insects faster than they can be eliminated, allowing toxins to move through the soil food web. In addition, contaminants may alter the structure and diversity of soil microbial communities, eliminate sensitive species and allowing resistant ones to dominate, which ultimately disrupts ecological balance and nutrient cycling processes (Innocent *et al.*, 2024). Most of the trace metals are also cause phytotoxic effects, inhibiting root development, nutrient uptake, and plant growth. Furthermore, toxic substances can modify soil chemical properties such as pH and nutrient availability, indirectly affecting soil fertility and biological activity (Kabata-Pendias, 2011). Together, these mechanisms reduce soil productivity, biodiversity, and ecosystem stability, highlighting the importance of monitoring and managing soil contamination (FAO, 2018).

### ***Effects of metal toxicity on plant health***

Heavy metal toxicity severely impairs plant health by disrupting essential physiological and biochemical processes, primarily through the overproduction of reactive oxygen species (ROS) such as superoxide, hydrogen peroxide, and hydroxyl radicals, which cause oxidative stress

leading to lipid peroxidation, protein oxidation, DNA damage, and cellular redox imbalance (Gill and Tuteja, 2010; Bhat *et al.*, 2025). Non-essential metals like Cd, Pb, As, and Hg, as well as excess levels of essential ones such as Cu and Zn, interfere with nutrient uptake and translocation by competing with ions like Fe, Zn, Ca, and Mg at transport sites, resulting in deficiencies that further compromise metabolism. Photosynthesis is particularly vulnerable, with metals inducing chlorophyll degradation (*via* enhanced chlorophyllase activity or  $Mg^{2+}$  replacement in the porphyrin ring), inhibiting electron transport in photosystems (especially PS II), reducing  $CO_2$  fixation, causing chloroplast disorganization, and triggering photo-oxidative damage, all of which manifest as chlorosis, reduced photosynthetic efficiency, and lowered biomass (Clemens, 2006). Enzyme activities are inhibited through direct binding to sulfhydryl groups or oxidative inactivation, affecting key processes like respiration, nitrogen assimilation, and antioxidant defences, while membrane permeability is altered, leading to electrolyte leakage, osmotic stress, and impaired water relations. Visible symptoms include stunted growth, shortened internodes, root browning or necrosis (often starting at tips), leaf rolling or wilting, interveinal chlorosis (especially in young leaves), necrotic spots, premature senescence, delayed flowering, reduced pollen viability, poor seed set, and overall yield losses ranging from 20–

**Table 2: Key effect of heavy metals on plant health** (Nagajyoti *et al.*, 2010)

Metal/Metalloid	Essential	Common Sources in Soil	Typical Toxic Threshold (approx., mg/kg soil)	Key Affected Components
Cadmium (Cd)	No	Fertilizers, sewage sludge, smelting	>1–3 (very low tolerance)	Plants, microbes, humans (highly bioaccumulative)
Lead (Pb)	No	Historical gasoline/paint, mining, batteries	>100–400	Plants (roots), soil fauna, neurological effects
Arsenic (As)	No (some plants)	Pesticides (historical), mining, irrigation water	>10–50	Plants (root damage), carcinogen
Mercury (Hg)	No	Industrial, coal burning, artisanal mining	>1–5	Microbes, bioaccumulation in food chain
Chromium (Cr)	No	Tanneries, electroplating	>50–100	Oxidative damage
Copper (Cu)	Yes	Fungicides, fertilizers, mining	>100–300 (essential→toxic)	Roots, algae, invertebrates
Zinc (Zn)	Yes	Fertilizers, galvanizing, sewage	>200–400	Plants (chlorosis at excess), microbes
Nickel (Ni)	Yes	Mining, batteries, fertilizers	>50–150	Allergic/toxic effects
Manganese (Mn)	Yes	Natural, some fertilizers	>1000–5000 (acid soils mobilize)	Rare toxicity, but excess in acidic soils
Iron (Fe)	Yes	Natural, but excess rare	Very high (toxicity uncommon)	Rare, mostly in flooded/acidic conditions

60% or more in contaminated soils (Saha *et al.*, 2017). These effects highlight a narrow tolerance window for metals, contrasting their essential micronutrient roles at low concentrations, and underscore the need for remediation in affected agricultural areas like those in Rajasthan where fertilizer and irrigation sources contribute to contamination (Dotaniya *et al.*, 2016b).


**Metal contamination in the food chain**

Heavy metals and metalloids enter and contaminate the food chain primarily through polluted soils, where toxic elements like Cd, Pb, As, Hg, Cr, and excess Cu or Zn accumulate from anthropogenic sources such as industrial effluents, mining, phosphate fertilizers, sewage irrigation, and atmospheric deposition (Sharma *et al.*, 2007). Plants absorb these metals *via* roots from contaminated soil or through foliar uptake from polluted water/air, leading to bioaccumulation in edible parts like leaves, grains, roots, and fruits—often exceeding safe limits in crops such as rice, wheat, vegetables (e.g., spinach, leafy greens), and fodder (Singh *et al.*, 2010). This process transfers metals to herbivores (mostly livestock consuming contaminated plants or fodder), where they further accumulate in tissues, milk, and meat, and eventually reach humans as top consumers through the diet. Key mechanisms include bioaccumulation (buildup within an organism over time) and, for persistent non-essential

metals like Cd, Pb, and Hg, biomagnification (increasing concentrations at higher trophic levels), amplifying exposure risks despite lower initial soil levels. In terrestrial systems, unlike aquatic food webs where biomagnification is pronounced (e.g., Hg in fish), terrestrial chains show more variable transfer, but significant risks persist *via* direct plant-to-human or plant-to-animal-to-human pathways (Chary *et al.*, 2008). Globally, 14–17% of cropland exceeds toxic metal thresholds, potentially affecting 0.9–1.4 billion people through contaminated food. In regions like Rajasthan elevated Cd, Pb, and other metals in vegetables irrigated with wastewater or grown near industrial sites often linked to industrial pollution and groundwater use (Meena *et al.*, 2020). Human health consequences arise from chronic ingestion, causing organ damage (kidneys, liver, brain), neurological disorders, reproductive issues, and increased cancer risk, particularly in vulnerable populations reliant on local produce. Mitigation requires soil testing, clean irrigation, phytoremediation, and regulatory limits on fertilizers/sludge to break contamination pathways and safeguard food safety.

**Phytoremediation mechanisms for heavy metal remediation**

Phytoremediation is an eco-friendly, cost-effective, and sustainable technique that harnesses plants and their associated rhizosphere microorganisms to remediate heavy metal-contaminated soils and water by removing, degrading, stabilizing, or transforming pollutants through several key mechanisms depicted in Figure 2 (Mishra *et al.*, 2017; Oubohssaine and Dahmani, 2024). The primary mechanisms include phytoextraction (or phytoaccumulation), where plants absorb metals from soil *via* roots, translocate them to above ground shoots/leaves (often storing them in vacuoles as metal-phytochelatin or metal-ligand complexes to avoid toxicity), and allow harvest of metal-laden biomass for removal—ideal for metals like Cd, Pb, As, Zn, and Cu, with hyperaccumulators (*Pteris vittata* for As, *Sedum alfredii* or *Brassica* species for Cd) accumulating concentrations far exceeding soil levels (Jakovljevi *et al.*, 2024). Phytostabilization immobilizes metals in the rhizosphere by root adsorption, precipitation,



**Fig. 1: Type of phytoremediation process and mechanism in plants (Mishra *et al.*, 2017)**

or complexation (*via* root exudates or mycorrhizal fungi), reducing bioavailability, leaching into groundwater, and entry into the food chain—suitable for Pb, Cr, and As in mining-impacted sites. Rhizofiltration (or phytofiltration) uses plant roots (especially aquatic/hydroponic species) to adsorb, absorb, or precipitate metals from contaminated water or saturated zones, effectively treating wastewater or groundwater (Surriya *et al.*, 2015). Phytovolatilization involves uptake, transformation (e.g., methylation or reduction), and release of volatile metal forms (like Hg or As) into the atmosphere *via* transpiration, though this transfers rather than fully removes the contaminant and is limited to volatile elements (Rani *et al.*, 2023). Additional supportive processes include rhizodegradation (microbial breakdown in the root zone) and phytodegradation (internal enzymatic transformation). These mechanisms rely on plant tolerance strategies such as chelation by phytochelatins/metallothioneins, vacuolar sequestration, antioxidant defense against oxidative stress, and enhanced metal transporters (Zunaidi *et al.*, 2024). In regions like Rajasthan, where heavy metal contamination from industrial effluents, fertilizers, and irrigation water affects soils, local tolerant species (e.g., maize, *Physalis angulata*, *Acalypha indica*, or wetland plants) show promise for phytostabilization or phytoextraction, often enhanced by chelators like EDTA or microbial inoculants to boost efficiency. Overall, phytoremediation offers a green alternative to conventional methods, though success depends on plant selection, soil conditions, metal type/speciation, and time required for significant clean up (Kumar *et al.*, 2024).

#### ***Role of mustard in phytoremediation***

Mustard plants, particularly Indian mustard (*Brassica juncea*), play a prominent role in phytoremediation as a fast-growing, high-biomass crop from the Brassicaceae family, widely recognized for its ability to tolerate and accumulate various heavy metals from contaminated soils (Singh *et al.*, 2024). It functions primarily through phytoextraction (or phytoaccumulation), where roots absorb metals from the soil, translocate them

efficiently to shoots and leaves (often with high translocation factors), and store them in aboveground tissues—facilitated by metal-binding compounds like phytochelatins, metallothioneins, glutathione, and organic acids that chelate and sequester metals in vacuoles to minimize toxicity (Yang *et al.*, 2017). This species excels at extracting non-essential metals such as Cd (often 3 times more than many other plants), Pb (up to significant reductions like 13–36% in one season), As, Hg, Cr, and Se, as well as excess essential metals like Zn, Cu, and Ni. It also supports phytostabilization by binding metals in the rhizosphere *via* root exudates and associated microbes, reducing their bioavailability and preventing leaching or food-chain entry (Yadav *et al.*, 2016). Singh *et al.* (2024) conducted an experiment to identify the role of cultivar in heavy metal uptake dynamics; suggested that *B. juncea* cv. Varuna and NRCHB 101 could be used for the phytoextraction of heavy metals and reducing their contamination in food chain, respectively in wastewater irrigated areas of peri-urban India.

It's advantages include rapid growth, large biomass production, short life cycle, tolerance to moderate contamination (high tolerance index >90% in many studies), and activation of robust antioxidant defenses (e.g., SOD, CAT, APX) to combat oxidative stress from ROS induced by metals (Kesawat *et al.*, 2023). Efficiency is often enhanced by agronomic interventions such as applying chelators like EDTA (which increases metal solubility and shoot accumulation dramatically), sulfur amendments (to acidify soil and boost Cd uptake), peat or organic matter (improving growth and translocation), co-cropping with other plants (e.g., maize for combined remediation and production), or even induced mutations (e.g., EMS-treated genotypes for super-hyperaccumulation) (Zia-ur-Rehman *et al.*, 2023). In contaminated regions like parts of Rajasthan (including Kota's industrial/agricultural areas with wastewater irrigation), *B. juncea* varieties show promise for remediating Cd, Pb, and other metals in soils, though harvested biomass must be disposed of safely to avoid re-contamination. Overall, Indian mustard stands out as one of the most studied and effective “green” tools for cost-effective, *in-situ*

remediation of multi-metal polluted sites, balancing ecological restoration with agricultural feasibility. The Brassicaceae family is renowned for containing a large number of metal hyperaccumulators. More than 100 species within this family are known to hyperaccumulate metals, particularly Ni and Zn, with fewer for Cd, As, Pb, and others (Rathore *et al.*, 2019). Indian mustard (*Brassica juncea*), heavy metal uptake from contaminated soil follows a multi-step process involving root absorption, symplastic/apoplastic pathways, translocation *via* the xylem, and partitioning into various plant parts, often enhanced for phytoremediation purposes. Metals primarily exist as cations (e.g.,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Cu}^{2+}$ ,  $\text{Ni}^{2+}$ ) or anions (e.g., As as arsenate) in the soil solution, with bioavailability influenced by soil pH (higher uptake at lower pH), organic matter, and chelators (Zia-ur-Rehman *et al.*, 2023). The process begins at the root level through active uptake *via* specialized plasma membrane transporters. *B. juncea* utilizes transporters from families such as ZIP (Zrt/Irt-like proteins, e.g., for Zn, Fe, Cd), IRT (iron-regulated transporters, which also handle Cd and other divalent metals), NRAMP (natural resistance-associated macrophage proteins, involved in Cd, Mn, Fe transport), and others like YSL (yellow stripe-like) for metal chelates. These transporters facilitate entry into root epidermal and cortical cells, often mimicking essential micronutrient pathways (Cd competes with Zn/Fe uptake). Some metals (e.g., Pb) primarily enter *via* the apoplast (cell wall space) before crossing the Casparian strip in the endodermis into the symplast for selective transport (Pasricha *et al.*, 2021).

Once inside root cells, metals bind to chelators like phytochelators (PCs), glutathione (GSH), or organic acids to prevent toxicity and enable safe movement. From roots, metals are loaded into the xylem for upward translocation to shoots, driven by transpiration pull. Translocation factors (TF = shoot concentration/root concentration) vary by metal: higher for mobile metals like Zn, Cd (often >1 in *B. juncea*), and lower for immobile ones like Pb (mostly retained in roots) (Dotaniya *et al.*, 2018a). In stems and leaves, metals are distributed *via* phloem to some extent, but xylem dominates for most. Accumulation

is highest in leaves and stems for phytoextraction, with sequestration in vacuoles *via* tonoplast transporters like HMA (heavy metal ATPases), ABC transporters, or MTP (metal tolerance proteins) to detoxify and store them, often as metal-PC or metal-organic acid complexes (Pasricha *et al.*, 2021). For non-essential metals like Cd and As, uptake can occur *via* essential nutrient transporters (e.g., As *via* phosphate transporters), while chelators like EDTA (applied externally) enhance solubility, form stable complexes, and boost root-to-shoot translocation dramatically (e.g., Pb-EDTA complexes). In *B. juncea*, kinetic models describe uptake as first-order (rate proportional to concentration gradient), with time-dependent accumulation peaking in roots early and shifting to shoots over growth stages (Almeaiweed *et al.*, 2025). Overall, *B. juncea*'s efficiency stems from high root surface area, robust transporter expression, effective chelation, and strong antioxidant defences, enabling significant accumulation (e.g., high Cd/Pb in shoots under enhanced conditions) while maintaining growth—making it ideal for phytoextraction in contaminated soils in most of the industrial/agricultural areas.

### ***Mechanism of heavy metal uptake from soil to mustard plant parts***


The uptake and translocation of heavy metals in Indian mustard *Brassica juncea* follow a well-characterized pathway from soil solution to various plant parts, involving root absorption, internal transport, and sequestration. Below is a textual representation of the key steps, followed by relevant diagrams from scientific sources for visual clarity.

- i. Soil solution and rhizosphere phase metals (e.g.,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$ ,  $\text{Zn}^{2+}$ ,  $\text{Cu}^{2+}$ ) exist as ions or complexes in soil. Root exudates, microbial activity (e.g., siderophores from PGPR), and amendments (organic acids, chelators like EDTA) increase bioavailability by solubilizing insoluble forms (*via* proton release lowering pH or forming soluble complexes) (Kalyvas *et al.*, 2018).
- ii. Root uptake (absorption into root cells) metals enter root epidermal/cortical cells mainly *via* plasma membrane transporters (ZIP family for Zn/Cd/Fe, NRAMP for Cd/Fe/Mn, IRT for Fe/

- Cd). Non-essential metals like Cd often “hitchhike” on essential nutrient transporters. Pb tends to follow the apoplastic pathway (cell walls) before crossing the Casparian strip into the symplast (Pasricha *et al.*, 2021).
- iii. Chelation and detoxification in roots inside cells, metals bind to ligands like phytochelatins (PCs), glutathione (GSH), or organic acids to prevent toxicity and enable safe movement.
  - iv. Xylem loading and translocation to shoots metals are loaded into the xylem (*via* transporters like HMA family) and transported upward by transpiration pull. Translocation factors are often high for Cd/Zn (shoot > root), lower for Pb (mostly retained in roots).
  - v. Distribution and sequestration in shoots in stems/leaves, metals move to phloem to some extent but primarily accumulate in leaves. Sequestration occurs in vacuoles *via* tonoplast transporters (e.g., MTP, ABC, HMA3), often as metal-PC/GSH complexes, detoxifying and storing them in low-metabolic-activity compartments (Ali and Ahirwar, 2025).

### ***Why mustard crop (Brassica juncea) is commonly using in phytoremediation***

Mustard, specifically Indian mustard (*Brassica juncea*), is not the only plant used in phytoremediation—many species like *Thlaspi*



**Fig. 2: Heavy metal dynamics in different plant parts** (Bhat *et al.*, 2025)

*caerulescens* (Alpine pennycress for Zn/Cd hyperaccumulation), *Pteris vittata* (Chinese brake fern for As), sunflowers, corn, barley, alfalfa, and others serve effectively depending on the metal, soil type, and site conditions (Sheng *et al.*, 2008). However, *B. juncea* is one of the most frequently studied, recommended, and applied crops in research and field trials worldwide due to a unique combination of practical advantages that make it highly suitable for phytoextraction and related mechanisms.

Key reasons for its prominence include:

- High biomass production — It grows rapidly with substantial aboveground biomass (leaves and stems), allowing removal of larger total quantities of metals per plant or per hectare compared to low-biomass hyperaccumulators (e.g., many wild Brassicaceae species are slow-growing and small). Average biomass production of the mustard varieties ranged 3-10.3 q/ha including the stalks. However, the biomass production largely depends on the crop climate, cultivar, soil, agronomic management and plant nutrient management (Priyanka *et al.*, 2022).
- Fast growth and short life cycle — mustard is a hardy crop and the initial growth is very fast under the proper management of plant nutrients. It reaches maturity in 3–4 months and quicker results than perennial or slower plants.
- Broad metal accumulation capacity — It effectively accumulates and translocates a wide range of heavy metals (Cd, Pb, As, Hg, Cr, Zn, Cu, Ni, Se) from roots to shoots, often with high translocation factors, making it versatile for multi-metal contaminated sites rather than specialists for one metal (Dotaniya *et al.*, 2024).
- Good tolerance to moderate contamination — It withstands metal stress without severe toxicity (high tolerance index >90% in many studies), maintains growth, and activates strong antioxidant defences (e.g., SOD, CAT, APX) to counter ROS damage (Saha *et al.*, 2017).
- Ease of cultivation and agronomic management — As an agricultural crop (grown for greens, oilseeds, or fodder in regions like India), it is

well-known, with available varieties, seeds, and farming practices; it responds well to enhancements like chelators (EDTA dramatically boosts uptake), sulfur amendments, organic matter, or microbial co-inoculation (Pasricha *et al.*, 2021).

- High research backing and proven field success — Extensive studies (e.g., EPA trials showing 13% Pb reduction in one season with EDTA) and genotype screenings (e.g., certain cultivars outperforming others for Cd/Pb) have established it as a reliable model plant, leading to more publications and practical adoption compared to less-studied species (Pasricha *et al.*, 2021).

However, in addition above facts not exclusive or always superior (e.g., true hyperaccumulators like *Noccaea caerulescens* concentrate metals at higher tissue levels without chelators, and grasses like barley may tolerate higher concentrations), *Brassica juncea*'s balance of biomass, speed, versatility, tolerance, and agricultural feasibility makes it a “go-to” choice for cost-effective, scalable phytoremediation—especially in developing regions or agricultural soils, where it can integrate into farming systems while remediating contaminants from fertilizers, wastewater, or industrial sources. Other plants are chosen when site-specific needs (e.g., extreme hyperaccumulation or wetland conditions) demand them.

### ***Management practices for improving phytoremediation efficiency of Indian mustard***

Several agronomic, chemical, biological, and biotechnological management practices can significantly enhance the efficiency of phytoremediation for heavy metal-contaminated soils, addressing limitations like low metal bioavailability, slow plant growth, limited biomass, and poor translocation to shoots. These practices boost metal uptake (especially in phytoextraction), improve plant tolerance to stress, increase biomass production, and accelerate overall contaminant removal while minimizing environmental risks.

- Chemical-assisted approaches (Chelate application) — Adding synthetic chelators like

EDTA (ethylenediaminetetraacetic acid) or natural ones like citric acid increases metal solubility and bioavailability by forming soluble metal-chelate complexes, facilitating root uptake and shoot translocation. EDTA is highly effective (often increasing phytoextraction by several folds, e.g., dramatic boosts in Cd, Pb, Zn, and Cu accumulation in plants like *Brassica juncea*), but high doses can harm plant growth or risk groundwater leaching due to persistence. Citric acid (and other organic acids) is biodegradable, eco-friendlier, promotes plant growth under stress, and enhances uptake without severe toxicity—often preferred for sustainable applications, though sometimes less potent than EDTA for certain metals.

- Soil amendments and organic matter addition — Incorporating biochar, compost, manure, humic substances, or crop residues (e.g., sugar beet residue) improves soil structure, nutrient availability, pH adjustment (acidification to mobilize metals in alkaline soils common in Rajasthan), and microbial activity. These enhance root development, reduce metal toxicity to plants, and increase phytoextraction or phytostabilization efficiency (Dotaniya *et al.*, 2022a).
- Microbe-assisted (PGPR and mycorrhizal fungi) — Inoculating with plant growth-promoting rhizobacteria (PGPR) (*Pseudomonas*, *Bacillus*, *Azospirillum*) or arbuscular mycorrhizal fungi (AMF) promotes plant growth *via* nitrogen fixation, siderophore production, hormone regulation, and antioxidant enhancement, while increasing metal mobilization and tolerance. This bio-augmentation often synergizes with plants like mustard, improving biomass and metal removal without chemical risks (Khawula *et al.*, 2025).
- Agronomic and crop management practices — Selecting high-biomass, tolerant varieties (e.g., specific genotypes of *Brassica juncea*); plant genotype and cultivar variation - Significant differences exist among *B. juncea* genotypes (e.g., IM-25, IM-13, IM-65 better for Cd; IM-79, IM-24 for Pb), influencing tolerance, translocation factor (TF), and accumulation

(some show hyperaccumulation traits for specific metals) (Gurajala *et al.*, 2019). In addition, optimizing planting density, intercropping, or co-cropping; applying fertilizers (N, P, S) judiciously to support growth without competing with metals; ensuring proper irrigation and weed control; and timing chelator application (e.g., late growth stage to avoid early toxicity). Multiple cropping cycles per year with fast-growing species accelerate remediation.

- Advanced biotechnological interventions — Genetic engineering (transgenic plants overexpressing metal transporters, chelators like metallothioneins, or stress-response genes) enhances tolerance and accumulation. Mutation breeding (e.g., EMS-induced hyperaccumulators in mustard) or nanoparticle additions can further boost efficiency, though these are more research-oriented (Kumar *et al.*, 2022).
- In regions like alkaline soils, wastewater irrigation, and fertilizer-derived Cd/Pb contamination are common—these practices are particularly relevant: combining *Brassica juncea* with low-dose citric acid or EDTA, organic amendments, and PGPR/AMF inoculation offers a balanced, locally adaptable strategy. Success depends on site-specific testing (soil pH, metal speciation, contamination levels), monitoring plant health, and safe disposal of metal-laden biomass (e.g., incineration or ashing) to prevent re-entry into the food chain (Dotaniya *et al.*, 2018b). Integrated approaches often yield the best results, transforming phytoremediation from a slow process into a more practical, sustainable tool for soil restoration.

## CONCLUSION

Metal is an essential for plant growth and when reaches to the threshold level, it will create problems for other nutrients during crop growth. However, discharge of metal contaminated effluent or waste generated material accumulated in natural resources and cause many challenges towards food chain contamination and ecosystem services. Assessment of the metal toxicity by computing the metal indices and execute the remediation strategies either

chemical or biological. Over the remediation projects stated that most of the chemical remediation concepts are needed huge investment, which is mostly not preferable in developing countries. However, Indian mustard (*Brassica juncea* L.) is a cost-effective agriculture crop, cultivated on a major part of Indian North states. Huge biomass potential has a curative window to reduce the metal toxicity in soil solution. The transfer factor of root to shoot or further in edible part like oil is almost negligible. In addition, management of Indian mustard by soil and agronomic practices will also improve the phytoremediation potential of major cultivars.

## REFERENCES

- Ali, S. and Ahirwar, R. (2025). Accumulation of toxic metals in vegetable crops and associated dietary exposure risks to human health. *Discover Environment*, 3, doi 10.1007/s44274-025-00373-w.
- Almeaiweed, N.H., Aloud, S.S., Alotaibi, K.D., Alotaibi, F. and Alshebel, B. (2025). Enhancing phytoremediation of heavy metal-contaminated aridic soil using olive mill wastewater, sulfur, and chelating agents. *Sustainability*, 17(8):3745. <https://doi.org/10.3390/su17083745>
- Arnon, D. I. and Stout, P. R. (1939). The essentiality of certain elements in minute quantity for plants with special reference to copper. *Plant Physiology*, 14(2): 371–375, <https://doi.org/10.1104/pp.14.2.371>
- Bhat, B.A., Rather, M.A., Bilal, T., Nazir, R., Qadir, R.U. and Mir, R.A. (2025). Plant hyperaccumulators: a state-of-the-art review on mechanism of heavy metal transport and sequestration. *Front. Plant Sci.*, 16:1631378. doi: 10.3389/fpls.2025.1631378
- Chary, N. S., Kamala, C. T. and Raj, D. S. S. (2008). Assessing risk of heavy metals from consuming food grown on sewage irrigated soils and food chain transfer. *Ecotoxicology and Environmental Safety*, 69(3): 513–524.
- Clemens, S. (2006). Toxic metal accumulation, responses to exposure and mechanisms of

- tolerance in plants. *Biochimie*, 88(11), 1707–1719. doi: 10.1007/s00425-021-03642-z
- Dehkordi, M. M., Nodeh, Z. P., Dehkordi, K. S., Khorjestan, R. R. and Ghaffarzadeh, M. (2024). Soil, air, and water pollution from mining and industrial activities: Sources of pollution, environmental impacts, and prevention and control methods. *Results Engineering*, 102729.
- Donald, L. S. (2003). *Environmental Soil Chemistry*. Academic Press, 350p.
- Dotaniya, M.L., Dotaniya, C.K., Kumar, K., Yadav, R.K., Doutaniya, R.K., Meena, H.M., Yadav, D.K., Shukla, S., Shirale, A.O. and Lata, M. (2024). Ecosystem services from rehabilitated waste dumpsites. In: V.K. Pandey (Ed.), *Biodiversity and Ecosystem Services on Post-Industrial Land Vol. 1*, John Wiley & Sons Ltd., ISBN 978-1-394-18738-6, Pp. 329-355.
- Dotaniya, M.L., Dotaniya, C.K., Yadav, D.K., Mahmoud, A.E.D., Doutaniya, R.K., Kumar, D., Kumar, A., Meena H.M., Verma R., Kuldeep, K. and Meena, A. (2023). Chromium dynamics in the soil-plant continuum. In: Kumar *et al.* (Eds), *chromium in plants and environment*. Springer Nature Switzerland doi.org/10.1007/978-3-031-44029-8
- Dotaniya, M.L., Meena, V.D., Dotaniya, C.K., Meena, M.D., Doutaniya, R.K., Verma, R., Sanwal, R.C., Parewa, H.P., Jatav, H.S., Meena, R., Sarkar, A. and Saha, J.K. (2022). Engineered biochar as adsorbent for removal of heavy metals from soil medium. In: S. Ramola, D. Mohan, O. Masek, A. Méndez, T. Tsubota (Eds), *Engineered Biochar*. Springer, Singapore. [https://doi.org/10.1007/978-981-19-2488-0\\_9](https://doi.org/10.1007/978-981-19-2488-0_9)
- Dotaniya, M.L., Meena, V.D., Rajendiran, S., Coumar, M.V., Saha, J.K., Kundu, S. and Patra, A.K. (2016). Geo-accumulation indices of heavy metals in soil and groundwater of Kanpur, India under long term irrigation of tannery effluent. *Bull Environ Conta Toxic* doi:10.1007/s00128-016-1983-4
- Dotaniya, M.L., Panwar, N.R., Meena, V.D., Regar, K.L., Lata, M. and Saha, J.K., (2018). Bioremediation of metal contaminated soils for sustainable crop production. In: V.S. Meena (Ed.), *Role of rhizospheric microbes in soil*, Springer International, Pp. 143-173.
- Dotaniya, M.L., Rajendiran, S., Dotaniya, C.K., Solanki, P., Meena, V.D., Saha, J.K. and Patra, A.K. (2018). Microbial assisted phytoremediation for heavy metal contaminated soils. In: V. Kumar, M. Kumar, R. (Eds) *Phytobiont and ecosystem restitution*. Springer, Pp. 295-317.
- Dotaniya, M.L., Rajendiran, S., Meena, V.D., Saha, J.K., Coumar, M.V., Kundu, S. and Patra, A.K. (2016). Influence of chromium contamination on carbon mineralization and enzymatic activities in Vertisol. *Agriculture Research*, doi: 10.1007/s40003-016-0242-6.
- Dotaniya, M.L., Rajendiran, S., Saurabh, K., Saha, J.K., Dotaniya, C.K. and Patra, A.K. (2022). Immobilization of chromium bioavailability through application of organic waste to Indian mustard (*Brassica juncea*) under chromium contaminated Indian soils. *Environment Monitoring and Assessment*, Doi 10.1007/s10661-022-10625-4.
- FAO (2018). *Food and Agriculture Organization. Soil Pollution: A Hidden Reality*. FAO, Rome.
- Gill, S. S. and Tuteja, N. (2010). Reactive oxygen species and antioxidant machinery in abiotic stress tolerance in crop plants. *Plant Physiology and Biochemistry*, 48(12): 909–930.
- Gupta, N., Khan, D. K. and Santra, S. C. (2012). Heavy metal accumulation in vegetables grown in sewage irrigated soils of Kolkata, India: A case study. *Ecotoxicology and Environmental Safety*, 94: 153–160.
- Gurajala, H.K, Cao, X., Tang, L., Ramesh, T. M., Lu, M. and Yang, X. (2019). Comparative assessment of Indian mustard (*Brassica juncea* L.) genotypes for phytoremediation of Cd and Pb contaminated soils.

- Environmental Pollution*, 254: Part B,113085.
- Innocent, M. O., Mustapha, A., Abdulsalam, M., Livinus, M. U., Samuel, J. O., Elelu, S.-A., Lateefat, S. O. and Muhammad, A. S. (2024). Soil microbes and soil contamination. In: J. C. Andrade, S. Mafla, E. Avila, J. Hernández (Eds.). *Soil Microbiome in Green Technology Sustainability*, Springer Nature, Pp. 3-35.
- Jakovljevi, K., Mišljenovi, T., van der Ent A, Baker, A.J.M., Invernón, V.R. and Echevarria, G. (2024) Mining the herbarium for hyperaccumulators: Discoveries of nickel and zinc (hyper)accumulation in the genus *Noccaea* (Brassicaceae) through X-ray fluorescence herbarium scanning. *Ecological Research*. <https://doi.org/10.1111/1440-1703.12448>
- Kabata-Pendias A. (2011). *Trace Elements in Soils and Plants* (4th ed.). CRC Press, 420p.
- Kalyvas, G., Tsitselis, G., Gasparatos, D. and Massas, I. (2018). Efficacy of EDTA and olive mill wastewater to enhance As, Pb, and Zn phytoextraction by *Pteris vittata* L. from a soil heavily polluted by mining activities. *Sustainability*, 10: 1962.
- Kaur, H. and Garg, N. (2021). Zinc toxicity in plants: A review. *Planta*, 253:1–28.
- Kesawat, M.S., Satheesh, N., Kherawat, B.S., Kumar, A., Kim, H.U., Chung, S.M. and Kumar, M. (2023). Regulation of reactive oxygen species during salt stress in plants and their crosstalk with other signaling molecules-current perspectives and future directions. *Plants (Basel)*,12(4):864.
- Khawula, S., Daniel, A.I., Nyawo, N., Ndlazi, K., Sibiya, S., Ntshalintshali, S., Nzuzwa, G., Gokul, A., Keyster, M., Klein, A., Niekerk, L. and Nkomo, M. (2025). Optimizing plant resilience with growth-promoting Rhizobacteria under abiotic and biotic stress conditions. *Plant Stress*, 17:100949.
- Kumar, A., Dadhwal, M., Mukherjee, G., Srivastava, A., Gupta, S. and Ahuja, V. (2024). Phytoremediation: sustainable approach for heavy metal pollution. *Scientifica* (Cairo), doi: 10.1155/2024/3909400.
- Kumar, K., and Shinde, A., Aeron, V., Verma, A. and Arif, N. (2022). Genetic engineering of plants for phytoremediation: advances and challenges. *Journal of Plant Biochemistry and Biotechnology*, doi 10.1007/s13562-022-00776-3.
- Meena, M.K., Singh, A.K., Prasad, L.K., Islam, A., Meena, M.D., Dotaniya, M.L., Singh, H.V., and Yadav, B.L. (2020). Impact of arsenic-polluted groundwater on soil and produce quality: a food chain study. *Environment Monitoring Assessment*, 192: 785.
- Minhas, P.S., Saha, J.K., Dotaniya, M.L., Saha, A. and Saha, M. (2021). Wastewater irrigation in India: Current status, impacts and response options. *Science of the Total Environment*, 808:1-17.
- Mishra, J, Singh, R. and Arora, N.K. (2017). Alleviation of Heavy Metal Stress in Plants and Remediation of Soil by Rhizosphere Microorganisms. *Front. Microbiol.*, 8:1706. doi: 10.3389/fmicb.2017.01706
- Nagajyoti, P. C., Lee, K. D. and Sreekanth, T. V. M. (2010). Heavy metals, occurrence and toxicity for plants: A review. *Environmental Chemistry Letters*, 8: 199–216.
- Ojuederie, O. B., Igwe, D. O. and Popoola, J. O. (2022). Transgenic plant-mediated phytoremediation: Applications, challenges, and prospects. *Assisted Phytoremediation*, 179–202.
- Oubohssaine, M. and Dahmani, I. (2024). Phytoremediation: Harnessing plant power and innovative technologies for effective soil remediation. *Plant Stress*,14:100578.
- Pasricha, S., Mathur, V., Garg, A., Lenka, S., Verma, K. and Agarwal, S. (2021). Molecular mechanisms underlying heavy metal uptake, translocation and tolerance in hyperaccumulators-an analysis: Heavy metal tolerance in hyperaccumulators. *Environmental Challenges*, 4:100197.
- Pepper I., Gerba, C.P. and Gentry, T. J. (2014). *Environmental Microbiology* (3rd ed.). Academic Press, 123p.
- Priyanka, Prakash, R., Ankus, Devi, S., Rajpaul and

- Singh, R. (2022). Physiological studies and nutrient accumulation in mustard varieties in response to fertilizer doses under saline condition in semi-arid region of northwestern Haryana. *Indian Journal of Experimental Biology*, 60: 628-634.
- Rajendiran, S., Singh, T.B., Saha, J.K., Coumar, J.K., Dotaniya, M.L., Kundu, S. and Patra, A.K. (2018). Spatial distribution and baseline concentration of heavy metals in swell-shrink soils of Madhya Pradesh, India. In: V. Singh, S. Yadav, R. Yadava (Eds), *Environmental Pollution. Water Science and Technology Library*, Springer, Pp. 135-145.
- Rani, P., Rose, P.K., Kidwai, M.K. and Meenakshi (2023), *Brassica Juncea L.: A Potential Crop for Phytoremediation of Various Heavy Metals*. In: R.P. Singh, P. Singh, A. Srivastava (Eds.), *Heavy metal toxicity: environmental concerns, remediation and opportunities*, Springer Nature, Singapore, Pp. 285-311.
- Rathore, S.S., Shekhawat, K. and Dass, A. *et al.* (2019). phytoremediation mechanism in Indian mustard (*Brassica juncea*) and its enhancement through agronomic interventions. *Proceedings of the National Academy of Sciences, India Section B: Biological Sciences*, 89: 419-427.
- Saha, J.K., Selladurai, R., Coumar, M.V., Dotaniya, M.L., Kundu, S. and Patra, A.K., (2017). Major inorganic pollutants affecting soil and crop quality. In: J.K. Saha, R. Selladurai, M.V. Coumar, M.L. Dotaniya, S. Kundu, A.K. Patra (Eds.), *Soil pollution-an emerging threat to agriculture*. Springer, Singapore, Pp. 75-104.
- Sharma, P. and Dubey, R. S. (2005). Lead toxicity in plants. *Brazilian Journal of Plant Physiology*, 17(1): 35-52.
- Sharma, R. K., Agrawal, M. and Marshall, F. M. (2007). Heavy metal contamination of soil and vegetables in suburban areas of Varanasi, India. *Ecotoxicology and Environmental Safety*, 66(2): 258-266.
- Sheng, X.F., Xia, J.J., Jiang, C.Y., He, L.Y. and Qian, M. (2008). Characterization of heavy metal-resistant endophytic bacteria from rape (*Brassica napus*) roots and their potential in promoting the growth and lead accumulation of rape. *Environmental Pollution*, 156:1164-1170.
- Shiyab, S., Chen, J., Han, F. X., Monts, D. L., Matta, F. B., Gu, M. Su, Y. (2009). Phytotoxicity of mercury in Indian mustard (*Brassica juncea* L.). *Ecotoxicology and Environmental Safety*, 72: 619-625. doi: 10.1016/j.ecoenv.2008.06.002
- Singh, A., Sharma, R. K., Agrawal, M. and Marshall, F. M. (2010). Risk assessment of heavy metal toxicity through contaminated vegetables from wastewater irrigated area of Varanasi, India. *Tropical Ecology*, 51(2): 375-387.
- Singh, R., Gautam, N., Mishra, A. and Gupta, R. (2011) Heavy metals and living systems: An overviews. *Indian Journal of Pharmacology*, 43 (3): 246-253.
- Singh, P.K., Yadav, J.S. and Kumar, I. *et al.* (2024). Screening of mustard cultivars for phytoremediation of heavy metals contamination in wastewater irrigated soil systems. *Environmental Monitoring and Assessment*, 196: 321 <https://doi.org/10.1007/s10661-024-12506-4>
- Sparks, D. L. (2003). *Environmental soil chemistry: An overview*. In *Environmental Soil Chemistry*, Elsevier, Pp. 1-42, <https://doi.org/10.1016/b978-012656446-4/50001-3>
- Surriya, O., Saleem, S.S., Waqar, K. and Kazi, A.G. (2015). Phytoremediation of Soils: Prospects and Challenges. In: K.R. Hakeem, M. Sabir, M. Ozturk, A.R. Mermut (Eds.), *Soil Remediation and Plants*, Academic Press, Pp. 1-36.
- Yadav, P., Kaur, R., Kohli, S. K., Sirhindi, G. and Bhardwaj, R. (2016). Castasterone assisted accumulation of polyphenols and antioxidant to increase tolerance of *B. Juncea* plants towards copper toxicity. *Cogent Food & Agriculture*, 2(1): 1276821.
- Yang, Y., Zhou, X., Tie, B., Peng, L., Li, H., Wang,

- K. and Zeng, Q. (2017). Comparison of three types of oil crop rotation systems for effective use and remediation of heavy metal contaminated agricultural soil. *Chemosphere*, 188: 148–156.
- Zia-ur-Rehman, M., Mubsher, A., Rizwan, M., Usman, M., Jafir, M., Umair, M., Alharby, H., Bamagoos, A., Alshamrani, R. and Ali, S. (2023). Effect of farmyard manure, elemental sulphur and EDTA on growth and phytoextraction of cadmium by spider plants (*Chlorophytum comosum* L.) under Cd stress. *Chemosphere*, 313: 1-11.
- Zunaidi, A.A., Lim, L.H. and Metali, F. (2024). Heavy metal tolerance and accumulation in the Brassica species (*Brassica chinensis* var. *parachinensis* and *Brassica rapa* L.): A pot experiment. *Heliyon* doi 10.1016/j.heliyon.2024.e29528.

*Received: March 17, 2025*

*Accepted: April 29, 2025*