

Print ISSN : 0972-8813
e-ISSN : 2582-2780

[Vol. 24(2) May-August 2026]

Pantnagar Journal of Research

(Formerly International Journal of Basic and
Applied Agricultural Research ISSN : 2349-8765)



G.B. Pant University of Agriculture & Technology
Pantnagar, U.S. Nagar; Uttarakhand, Website : gbpuat.res.in/PJR



ADVISORY BOARD

Patron

Prof. Shivendra Kumar Kashyap, Ph.D., Vice-Chancellor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Members

Prof. S. K. Verma, Ph.D., Director Research, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Jitendra Kwatra, Ph.D., Director, Extension Education, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. S.S. Gupta, Ph.D., Dean, College of Technology, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. D. Kumar, Ph.D., Dean, College of Veterinary & Animal Sciences, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Alka Goel, Ph.D., Dean, College of Community Science, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. R.S. Jadoun, Ph.D., Dean, College of Agribusiness Management, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Lokesh Varshney, Ph.D., Dean, College of Post Graduate Studies, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Avdhesh Kumar, Ph.D., Dean, College of Fisheries, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Anil Shukla, Ph.D., Dean, College of Agriculture, G.B. Pant University of Agri. & Tech., Pantnagar, India

Prof. Ramesh Chandra Srivastava, Ph.D., Dean, College of Basic Sciences & Humanities, G.B.P.U.A.T., Pantnagar, India

EDITORIAL BOARD

Members

A.K. Misra, Ph.D., Ex-Chairman, Agricultural Scientists Recruitment Board, Krishi Anusandhan Bhavan I, New Delhi, India & Ex-Vice Chancellor, G.B. Pant University of Agriculture & Technology, Pantnagar

Anand Shukla, Director, Reefberry Foodex Pvt. Ltd., Veraval, Gujarat, India

Anil Kumar, Ph.D., Director, Education, Rani Lakshmi Bai Central Agricultural University, Jhansi, India

Ashok K. Mishra, Ph.D., Kemper and Ethel Marley Foundation Chair, W P Carey Business School, Arizona State University, U.S.A

Binod Kumar Kanaujia, Ph.D., Professor, School of Computational and Integrative Sciences, Jawahar Lal Nehru University, New Delhi, India

D. Ratna Kumari, Ph.D., Associate Dean, College of Community / Home Science, PJTSAU, Hyderabad, India

Deepak Pant, Ph.D., Separation and Conversion Technology, Flemish Institute for Technological Research (VITO), Belgium

Desirazu N. Rao, Ph.D., Honorary Professor, Department of Biochemistry, Indian Institute of Science, Bangalore, India

G.K. Garg, Ph.D., Ex-Dean, College of Basic Sciences & Humanities, G.B. Pant University of Agric. & Tech., Pantnagar, India

Humnath Bhandari, Ph.D., IRRI Representative for Bangladesh, Agricultural Economist, Agrifood Policy Platform, Philippines

Indu S. Sawant, Ph.D., Principal Scientist, ICAR National Research Centre for Grapes, Pune, India

Kuldeep Singh, Ph.D., Director, ICAR - National Bureau of Plant Genetic Resources, New Delhi, India

Muneshwar Singh, Ph.D., Ex-Project Coordinator AICRP- LTFE, ICAR, Indian Institute of Soil Science, Bhopal, India

Omkar, Ph.D., Professor (Retd.), Department of Zoology, University of Lucknow, India

P.C. Srivastav, Ph.D., Professor (Retd.), Department of Soil Science, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Prashant Srivastava, Ph.D., Soil Contaminant Chemist, CSIRO, Australia

Puneet Srivastava, Ph.D., Director, Water Resources Center, Butler-Cunningham Eminent Scholar, Professor, Biosystems Engineering, Auburn University, United States

R.K. Singh, Ph.D., Ex-Director & Vice Chancellor, ICAR-Indian Veterinary Research Institute, Izatnagar, U.P., India

Ramesh Kanwar, Ph.D., Charles F. Curtiss Distinguished Professor of Water Resources Engineering, Iowa State University, U.S.A.

S.N. Maurya, Ph.D., Professor (Retired), Department of Gynaecology & Obstetrics, G.B. Pant University of Agric. & Tech., Pantnagar, India

Sham S. Goyal, Ph.D., Professor Emeritus, Faculty of Agriculture and Environmental Sciences, University of California, Davis, U.S.A.

Umesh Varshney, Ph.D., Honorary Professor, Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, India

V.D. Sharma, Ph.D., Dean Life Sciences, SAI Group of Institutions, Dehradun, India

V.K. Singh, Ph.D., Director, ICAR-Central Research Institute for Dryland Agriculture, Hyderabad, India

Vijay P. Singh, Ph.D., Distinguished Professor, Caroline and William N. Lehrer Distinguished Chair in Water Engineering, Department of Biological and Agricultural Engineering, Texas A & M University, U.S.A.

Nikitin Georgy, Ph.D., Vice-Rector cum Professor St. Petersburg State University of Veterinary Medicine St. Petersburg, Russia

Sanjay Jambhulkar, Ph.D., Ex-Scientific Officer 'H' Head Mutation Breeding Section Bhabha Atomic Research Centre, Mumbai Professor, Homi Bhabha National Institute, Mumbai

Vijay Kubsad, Ph.D., Adjunct Professor Washington State University USA

Kiran Tota-Maharaj, Ph.D., Professor Royal Agricultural University (RAU) Cirencester, Gloucestershire GL76JS England, UK

Ravindra P. Veeranna, Ph.D., Dean, Xavier University, College of Veterinary Sciences, Aruba

Javed Akhtar, Ph.D., University of Technology and Applied Sciences, Muscat

Basant Maheshwari, Ph.D., Western Sydney University, Australia

Vijay P. Singh, Ph.D. Professor Emeritus, Texas A&M University College station, Texas, USA

Sanjay Sharma, Ph.D. Director, Uttarakhand Council of Biotechnology

Krishna Nemali, Ph.D. Controlled Environment Agriculture Extension Specialist

Editor-in-Chief

K.P. Raverkar, Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Managing Editor

S.K. Guru, Ph.D., Professor, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Assistant Managing Editor

Jyotsna Yadav, Ph.D., Research Editor, Directorate of Research, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Technical Manager

S.D. Samantaray, Ph.D., Professor & Head, Department of Computer Engineering, G.B. Pant University of Agriculture and Technology, Pantnagar, India

Development

Dr. S.D. Samantaray, Professor & Head

Brijesh Dumka, Developer & Programmer

PANTNAGAR JOURNAL OF RESEARCH

Vol. 24(2)

May-August, 2026

CONTENTS

- AI and geospatial innovations for Indian agriculture: Applications, policy support and emerging research frontiers (Review)** 233-247
ABHISHEK DANODIA, SURESH KUMAR and AJEET SINGH NAIN
- Optimizing nitrogen use efficiency and grain quality in basmati rice through Integrated weed management: Insights into nutrient partitioning and Crop-weed competition** 248-255
ROMAN NISSAR, MANZOOR A GANAI, MEHRAJ UD DIN SOFI, ASHAQ HUSSAIN and F. A. BAHAR
- Performance evaluation of brown top millet (*Brachiaria ramosa* L.) genotypes under rainfed upland ecology of the Bastar plateau of Chhattisgarh** 256-266
PRAFULL KUMAR
- Effect of pre-harvest sprays of nanofertilizers on fruit quality in litchi (*Litchi chinensis* Sonn.) cv. Rose Scented** 267-274
NITIN PANWAR, RATNA RAI and RAJEEW KUMAR
- Ethrel induced ripening optimally modulates physicochemical quality, antioxidant properties and shelf life of sapota (*Manilkara zapota* L.) cv. Cricket Ball** 275-285
ASHOK KUMAR SINGH, KIRTI CHETRI, RISHABH RAJ, S. K. GURU and N. C. SHAHI
- Influence of drip irrigation and mulching practices on fruit set, yield and water use efficiency of litchi cv. rose scented in the Tarai region of Uttarakhand** 286-292
SHANKAR LAL, N. SINGH, N.K. BIND, PRERNA PRACHI, RAJSHREE, RIYA THAPA and AKHIL RUHELA
- Spatial distribution and survey of soybean anthracnose in the Kumaon region of Uttarakhand, India** 293-300
MANISHA DEV and K.P. SINGH
- Dimensionality reduction of part milk yield in crossbred cattle using Principal Component Analysis** 301-307
SIMRAN KAUR, ASHIS KUMAR GHOSH, OLYMPICA SARMA and RAVINDER SINGH BARWAL
- Comparative histomorphological and histochemical studies on the abomasum of adult Bakerwali goats, non-descript goats and non-descript sheep of the Jammu region** 308-318
JASVINDER SINGH SASAN, SHALINI SURI and KAMAL SARMA
- Gross morphology and biometry on the abomasum of adult Bakerwali goat, non-descript goats and non-descript sheep of Jammu region: A comparative study** 319-325
JASVINDER SINGH SASAN, SHALINI SURI and KAMAL SARMA

Arterial blood gas analysis in bovine pneumonia: Comparative study of healthy and diseased cattle ASMITA NARANG and CHARANJIT SINGH	326-331
Assessment of commonly used insecticide residues using gas chromatography in buffalo liver samples collected from North India S. K. JAIN, VINOD KUMAR, VINAY KANT, PREETI BAGRI, ARCHANA LOHIYA, GAURI CHANDRATRE and GAURAV GUPTA	332-339
<i>In silico</i> molecular docking analysis of squalene with major ageing-related molecular targets: matrix metalloproteinase-1, matrix metalloproteinase-9, cyclooxygenase-2, Tumor Necrosis Factor-α, hyaluronidase-1 and B-cell lymphoma-2 PAWANESH TAMTA, KULDIP CHANDRA VERMA and APARNA DIXIT	340-352
Characterization of agro-waste (<i>Parthenium hysterophorus</i>) derived Carbon Dots and to assess their potential as a sustainable adsorbent NISHTHA NAUDIYAL and R.K. SRIVASTAVA	353-360
Assessment of the antibacterial efficacy of endophytic bacteria in combination with neem leaf powder against bacterial pathogens SHIVANI SINGH, AJAY VEER SINGH and VELLAICHAMY MAGESHWARAN	361-370
Exploration of endophytic bacterial isolates from Kasni leaves (<i>Cichorium intybus</i>) for Phyto-hormone and Phyto-stimulating metabolite production VARSHA MISHRA, SEEMA BISHT, DIVYA PANT, ANUPAMA PANDEY and LAKSHMI TEWARI	371-382
Mitigation of dexamethasone-induced cardiac oxidative damage by <i>Moringa oleifera</i> and Vitamin E-Selenium: A biochemical study in broilers LATA PALIWAL, SONU AMBWANI and TANUJ KUMAR AMBWANI	383-395
Adaptive Neural Networks for dynamic prioritization in autonomous vehicles for road users MUMUKSHU BHATT, SUBODH PRASAD, RAJESH SINGH, ASHOK KUMAR and POOJA TAMTA	396-403
Assessing the impact of Geopolitical Disturbances on India's agricultural trade performance JYOTI, SUMAN GHALAWAT, MEGHA GOYAL, SUNIL KUMAR, AMITA GIRDHAR and SUBODH AGARWAL	404-409
Black soybean (<i>Glycine max</i> (L.) Merrill): Nutritional and functional analysis of whole seed, dehusked seed and husk fractions RITA SINGH RAGHUVANSHI, GUNJAN TEWARI, AABHA RAWAT and APURVA	410-421
<i>In silico</i> bioprospecting of bioactive phytochemicals with potential protective effects against exercise-induced muscle stiffness NAGARJUN G. S., PAL MURUGAN M., KRISHNANDU MONDAL, PADMASHREE A., WADIKAR D. D. and KUMAR R.	422-433

Ethrel induced ripening optimally modulates physicochemical quality, antioxidant properties and shelf life of sapota (*Manilkara zapota* L.) cv. Cricket Ball

ASHOK KUMAR SINGH^{1*}, KIRTI CHETRI¹, RISHABH RAJ¹, S. K. GURU² and N. C. SHAHI³

¹Department of Horticulture, College of Agriculture, ²Department of Plant Physiology, College of Basic Sciences and Humanities, G.B. Pant University of Agriculture and Technology, Pantnagar- 263145 (U. S. Nagar, Uttarakhand), ³Department of Processing and Food Engineering, MCAET, ANDUAT, Ambedkar Nagar-224122

*Corresponding author's email id: drashokkumar71@gmail.com

<https://doi.org/10.67063/pjr.vol.24.issue.02.05>

ABSTRACT: The present study was carried out during 2023–24 at GBPUAT, Pantnagar, Uttarakhand. The study was conducted to evaluate the effect of ethrel on ripening, fruit quality, antioxidant properties and shelf-life of sapota fruits. The experiment was conducted in a factorial Completely Randomized Design (CRD) with seven treatments and three replications. Mature fruits of sapota cv. Cricket Ball were treated with different concentrations of ethrel (500, 1000, 1500, 2000 and 2500 ppm), calcium carbide @ 10 g kg⁻¹ fruits, untreated control and stored under ambient conditions. Observations were recorded at 3, 6, 9 and 12 days after treatment. Significant differences were observed among treatments for physiological loss in weight, firmness, total soluble solids, total sugars, ascorbic acid, total phenolic content, total antioxidant activity and shelf-life. Fruits treated with Ethrel @ 1000 ppm showed more uniform ripening, higher TSS and total sugars, acceptable firmness, better antioxidant retention. The higher shelf-life was recorded with ethrel @ 500 ppm (10.50 days), which was statistically at par with ethrel @ 1000 ppm (10.33 days). However, CaC₂ treated fruits showed rapid deterioration, poor quality retention and lowest shelf-life. Overall, ethrel @ 1000 ppm provided a favourable balance between ripening uniformity, TSS and sugar accumulation, antioxidant retention and acceptable shelf-life under the ambient storage conditions of the study.

Keywords: Artificial ripening, CaC₂, Ethrel, Fruit quality, Sapota, Shelf-life

Sapota (*Manilkara zapota* L), popularly known as chikoo or sapodilla belongs to Sapotaceae family, is an important tropical evergreen fruit. It was originated from Southern Mexico and Central America and widely cultivated in tropical and subtropical regions of the world (Baskar *et al.*, 2020). India is one of the largest producers of sapota globally, covering an area of about 69.85 thousand hectares and production exceeding 861 thousand metric tonnes annually (MoA&FW, 2025). Gujarat, Karnataka, Maharashtra Andhra Pradesh, Tamil Nadu and Telangana are major sapota growing states of India. Important cultivars like Cricket Ball, Kalipatti, Jhumari, Pala and Kirthibarthi are widely cultivated because of superior fruit quality and market demand (Rekha, 2024). They are valued for their superior taste, pleasing aroma with high nutritional composition. It holds good amounts of sugars, proteins, minerals, carotenoids, phenolic compounds and antioxidants, which contribute significantly to their nutraceutical and health

promoting properties (Kaur *et al.*, 2020). Sapota holds an important rank among fruit crops in local as well as international markets due to its nutritional wealth and consumer suitability (Srivastava *et al.*, 2017).

It holds immense commercial importance but sapota fruits are highly perishable due to its climacteric nature and fast ripening behaviour after harvest. It exhibits quick climacteric respiratory pattern with high ethylene production during ripening (Baidya *et al.*, 2020). Due to which it suffers from fast ripening after harvest resulting in softening, high physiological loss in weight, biochemical alteration, microbial infection and accelerated senescence which reduces its shelf life and market suitability (Madani *et al.*, 2018). Sapota fruits were generally stored for 5-6 days after harvest under ambient conditions (Kaur *et al.*, 2020). However, considerable postharvest losses ranging between 20–35% have been reported during post-harvest

handling and marketing (Baidya *et al.*, 2020).

In commercial fruit markets of developing countries, calcium carbide (CaC_2) is still used as a low-cost ripening method despite its ban and associated health issue. It produces acetylene gas in the presence of moisture, which replicates the action of ethylene and induces rapid ripening (Vidhya *et al.*, 2025). Fruits ripened with CaC_2 shows reduced flavour, lower nutritional quality, extreme softening, quick senescence and reduced storage life (Desai *et al.*, 2024). Furthermore, CaC_2 available in local market contains harmful impurities like arsenic and phosphorus compounds that carries serious health risks to human health (Oladipupo *et al.*, 2022). There have been several studies on artificial ripening on climacteric fruits like mango, banana, papaya but with limited information available on sapota with various concentration of ethrel and CaC_2 on ripening, storage life and fruit quality.

Therefore, the present study was conducted to study the effect of different ethrel concentrations on ripening, fruit quality and shelf life of sapota cv. Cricket Ball under ambient conditions. The study also aimed to identify a suitable ethrel concentration for uniform ripening, better fruit quality and longer shelf-life and to compare its effects with calcium carbide ripening.

MATERIAL AND METHODS

Experimental Site and Plant Material

The study was conducted during 2023-24 at the Horticulture Research Centre, Patharchatta and the Department of Horticulture, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar, Uttarakhand, India. During May–June 2024, ambient temperature ranged from 34.1–42.6 °C (maximum) and 17.8–27.9 °C (minimum), while relative humidity ranged from 51.4–80.7% (maximum) and 14.1–65.0% (minimum). Physiologically mature fruits of sapota (*Manilkara zapota* L.) cv. Cricket Ball were harvested from healthy and uniformly managed trees. Fruits were selected based on uniform size, shape and maturity characteristics.

Experimental Design and Treatments

The experiment was conducted under ambient laboratory conditions in a two-factorial Completely Randomized Design (CRD) with three replications. Seven post-harvest ripening treatments were evaluated namely ethrel at 500 ppm (T_1), 1000 ppm (T_2), 1500 ppm (T_3), 2000 ppm (T_4) and 2500 ppm (T_5), calcium carbide at 10 g kg^{-1} fruit (T_6) and control (T_7). Each treatment in each replication consisted of 12 fruits, with three fruits used for evaluation at each storage interval (3rd, 6th, 9th and 12th day). Different fruits were used at each storage interval.

Fruits treated with different ethrel treatments were dipped in freshly prepared ethrel solutions for 5 min, air dried for 1 hour, packed in cardboard boxes and stored under ambient conditions. Calcium carbide treated fruits were packed in cardboard boxes with calcium carbide @ 10 g kg^{-1} fruit wrapped in newspaper and placed separately from the fruits to avoid direct contact, whereas control fruits were stored without treatment.

Determination of Quality Parameters

Physiological Loss in Weight (PLW) (%)

PLW was determined using an electronic balance and expressed as percentage loss relative to initial fruit weight using the formula:

$$\text{PLW (\%)} = \frac{\text{Initial fruit weight} - \text{Fruit weight on observation day}}{\text{Initial fruit weight}} \times 100$$

Fruit Firmness (N)

Fruit firmness was measured using a Texture Analyzer (TA. XT Plus, Stable Micro Systems, UK) fitted with a 2 mm probe. The force required to penetrate the fruit pulp to a depth of 10 mm was recorded and expressed in Newton (N).

Total Soluble Solids (TSS) (°Brix)

Total soluble solids were estimated using a hand refractometer and expressed as °Brix. Fresh fruit juice was placed on the prism surface and readings were recorded directly.

Ascorbic Acid (mg/100 g)

Ascorbic acid content was estimated by the 2,6-dichlorophenol indophenol dye titration method and calculated using the following formula:

$$\text{Ascorbic acid (mg/100 g)} = \frac{\text{Titre value} \times \text{Dye factor} \times \text{Volume made up}}{\text{Aliquot taken} \times \text{Weight of sample}} \times 100$$

Total Sugars (%)

Total sugars were estimated by the Lane and Eynon method using Fehling's solution and methylene blue indicator. Total sugars were calculated using the formula:

$$\text{Total sugars (\%)} = \frac{\text{Factor for Fehling's solution} \times \text{Dilution} \times 100}{\text{Titre value} \times \text{Weight of sample}}$$

Total Phenolic Content (mg GAE/100 g FW)

Total phenolic content was determined by the Folin–Ciocalteu colorimetric method. Absorbance was measured at 760 nm using a UV–Visible spectrophotometer and results were expressed as mg gallic acid equivalents (GAE) per 100 g fresh weight using a gallic acid standard curve.

Total Antioxidant Activity (%)

Total antioxidant activity was estimated using the DPPH free radical scavenging assay and calculated using the following formula:

$$\text{DPPH scavenging activity (\%)} = \left(1 - \frac{\text{Absorbance of sample}}{\text{Absorbance of control}}\right) \times 100$$

Shelf-life (Days)

Shelf life was determined based on visual assessment of marketability, including development of shrivelling, excessive softening and decay symptoms. It was expressed as the number of days fruits remained acceptable under ambient storage conditions.

Statistical Analysis

The experimental data were subjected to analysis of variance (ANOVA) for a 7×4 factorial Completely Randomized Design (CRD), considering ripening treatment, storage interval and their interaction ($T \times I$). The significance of treatment, storage interval and $T \times I$ interaction was tested at the 5% probability level. Treatment means were compared using the critical difference (CD) at $P = 0.05$ following Snedecor and Cochran (1987).

RESULTS AND DISCUSSION

Physiological Loss in Weight (%)

Physiological loss in weight (PLW) of sapota cv. Cricket Ball was significantly influenced by ripening treatments, storage duration and their interaction

(Fig. 1). PLW increased progressively with increasing concentrations of ethrel and advancement of storage period. Among the treatments, maximum PLW (16.70%) was recorded in calcium carbide treated fruits (T_6), followed by ethrel @ 2500 ppm (T_5) with 14.34%, whereas minimum PLW (6.13%) was observed in untreated control fruits (T_7). PLW increased significantly from the 3rd day (5.37%) to the 12th day (18.51%) across storage periods. Interaction effects were also significant, with the highest PLW (26.03%) recorded in calcium carbide treated fruits on the 12th day of storage, while the lowest PLW (1.85%) was observed in control fruits on the 3rd day. Lower ethrel concentrations, particularly 500 ppm (T_1) and 1000 ppm (T_2), maintained comparatively lower PLW throughout storage. The continuous increase in PLW during storage because of higher respiration, transpiration losses and fast consumption of stored metabolites during ripening (Rana *et al.*, 2021). Application of exogenous ethylene from ethrel hastens respiratory action, permeability of membrane and moisture loss which leads to higher fruit weight reduction (Tipu and Sherif, 2024). Fruit treated with CaC_2 displayed higher PLW due to acetylene induced quick ripening, tissue softening and accelerated desiccation (Vidhya *et al.*, 2025). Artificial ripening agents significantly enhanced metabolic activity and postharvest moisture loss as reported by Palpandian *et al.* (2019) in mango and Muthal *et al.* (2019) in banana.

Fruit Firmness (N)

Fruit firmness of sapota cv. Cricket Ball was significantly influenced by ripening treatments,





Fig. 1: Effect of post-harvest ethrel treatments and storage duration on physiological loss in weight (PLW, %) of sapota cv. Cricket Ball during ripening

Table 1: Effect of post-harvest ethrel treatments on total soluble solids (TSS), ascorbic acid and total sugar of sapota cv. Cricket Ball during ripening at different storage interval

Treatments	TSS (°Brix)				Ascorbic Acid (mg/100 g)				Total Sugars (%)			
	3 rd day	6 th day	9 th day	12 th day	3 rd day	6 th day	9 th day	12 th day	3 rd day	6 th day	9 th day	12 th day
T ₁ : Ethrel @ 500 ppm	16.57	18.93	21.23	18.03	22.30	18.36	15.98	11.76	11.02	13.19	15.59	13.36
T ₂ : Ethrel @ 1000 ppm	17.00	19.40	22.03	19.03	22.00	17.94	14.90	11.44	11.52	13.72	15.91	13.96
T ₃ : Ethrel @ 1500 ppm	17.17	20.63	19.03	16.37	21.83	16.71	14.10	10.13	11.83	14.64	12.75	12.01
T ₄ : Ethrel @ 2000 ppm	17.43	21.77	18.50	15.17	20.84	15.64	13.94	9.80	12.22	15.32	12.11	11.15
T ₅ : Ethrel @ 2500 ppm	17.80	21.97	18.00	14.83	20.00	14.87	12.06	9.15	12.54	15.76	11.85	10.20
T ₆ : CaC ₂ @ 10 g/kg fruit	18.93	17.50	15.57	13.20	18.50	12.93	10.26	6.21	13.88	12.95	10.85	9.39
T ₇ : Control	13.70	16.50	17.67	15.20	24.17	19.32	16.14	14.23	9.68	11.47	12.23	11.49
Mean	16.94	19.53	18.86	15.98	21.38	16.54	13.91	10.39	11.82	13.86	13.04	11.65
Parameters	Treatments (T)				Storage Intervals (I)				T × I			
TSS (°Brix)	0.50				0.38				0.99			
Ascorbic Acid (mg/100 g)	0.43				0.32				0.85			
Total Sugars (%)	0.40				0.30				0.79			

storage duration and their interaction (Fig. 2). Untreated control fruits (T₇) recorded the highest firmness (4.99 N), whereas CaC₂ treated fruits (T₆) exhibited the lowest firmness (1.45 N), followed by ethrel @ 2500 ppm (T₅) with 1.72 N among the treatments. Fruit firmness decreased significantly with advancement of storage period from 5.56 N on the 3rd day to 0.77 N on the 12th day, irrespective of treatments. The interaction effect was also significant, with maximum firmness (8.22 N) observed in control fruits on the 3rd day, while minimum firmness (0.27 N) was recorded in CaC₂ treated fruits on the 12th day, which was statistically at par with ethrel @ 2500 ppm. A rapid decline in firmness was observed in fruits treated with higher ethrel concentrations and CaC₂, indicating accelerated ripening and tissue softening (Vidhya *et al.*, 2025). More than 80% reduction in firmness occurred within six days in T₄, T₅ and T₆ treatments,

**Fig. 2: Effect of post-harvest ethrel treatments and storage duration on fruit firmness (N) of sapota cv. Cricket Ball during ripening**

coinciding with the climacteric rise in respiration. Softening of fruits due to higher degradation of cell wall polysaccharides, conversion of protopectin into soluble pectin and increased activities of cell wall degrading enzymes such as polygalacturonase, pectin methyl esterase and α -galactosidase (Liu *et al.*, 2026). Higher ethrel concentrations intensified enzymatic hydrolysis of structural carbohydrates, results in rapid softening of tissue and over ripening (Xia *et al.*, 2024). Similar findings were reported by Padmaja *et al.* (2015), Menezes and Athmaselvi (2016) and Vidhya *et al.* (2017) in sapota fruits.

Total Soluble Solids (°Brix)

Total soluble solids (TSS) of sapota cv. Cricket Ball was significantly influenced by ripening treatments, storage duration and their interaction (Table 1). Among the treatments, fruits treated with ethrel @ 1000 ppm (T₂) recorded the higher TSS (19.37 °Brix), followed by ethrel @ 500 ppm (T₁) with 18.69 °Brix, although control fruits (T₇) showed the lowest TSS (15.77 °Brix). Storage period significantly affected TSS, which increased from 16.94 °Brix on the 3rd day to 19.53 °Brix on the 6th day, followed by a gradual decline to 15.98 °Brix on the 12th day of storage. The interaction effect was also significant, with maximum TSS (22.03 °Brix) observed in fruits treated with ethrel @ 1000 ppm on the 9th day of storage. Minimum TSS (13.20 °Brix) was recorded in CaC₂ treated fruits on the 12th day. The increase in TSS during ripening might be

due to enzymatic hydrolysis of starch, pectic substances and complex carbohydrates into soluble sugars, whereas the subsequent decline could be associated with utilization of sugars during respiration (Yu *et al.*, 2022). Fruits treated with ethrel showed higher TSS due to enhanced ethylene facilitated activation of hydrolytic enzymes responsible for rapid carbohydrate conversion (Tipu and Sherif, 2024). Optimum concentrations of ethrel maintained higher TSS for a extended period, demonstrating balanced ripening and delayed senescence (Ghimire *et al.*, 2021). Similar findings were reported by Bons *et al.* (2016) in sapota, Muthal *et al.* (2019) in banana and Meiyarasan *et al.* (2025) in mango.

Ascorbic Acid (mg/100 g)

Ripening treatments, storage duration, and their interaction significantly influenced the ascorbic acid content of sapota cv. Cricket Ball (Table 1). Among the treatments, untreated control fruits (T_7) recorded the higher ascorbic acid content (18.46 mg/100 g), whereas CaC_2 treated fruits (T_6) showed the minimum value (11.97 mg/100 g). A significant decline in ascorbic acid was observed with advancement of storage period, decreasing from 21.38 mg/100 g on the 3rd day to 10.39 mg/100 g on the 12th day of storage. Interaction effects were also significant, with maximum ascorbic acid (24.17 mg/100 g) observed in control fruits on the 3rd day and minimum (6.21 mg/100 g) in CaC_2 treated fruits on the 12th day. Among ethrel treatments, fruits treated with ethrel @ 500 ppm maintained relatively higher ascorbic acid content throughout storage period. The gradual decline in ascorbic acid during ripening and storage due to greater oxidative breakdown and rapid consumption of antioxidants during climacteric respiration (Pareek, 2016). Fruits treated with ethrel showed earlier exhaustion of ascorbic acid due to accelerated ethylene mediated ripening and increased activity of oxidative enzymes responsible for conversion of ascorbic acid into dehydroascorbic acid (Asrey *et al.*, 2023). Lower retention of ascorbic acid in CaC_2 treated fruits because of acetylene induced metabolic stress and rapid senescence (Vidhya *et al.*, 2025). Similar findings were reported by Tsomu *et al.* (2015) in sapota, Adeyemi *et al.*

(2018) in banana, pawpaw and mango and Orisa and Usoroh (2020) in pawpaw fruits.

Total Sugars (%)

A significant effect of ripening treatments, storage period, and their interaction on the total sugars content of sapota cv. Cricket Ball was observed (Table 1). Among the treatments, fruits treated with ethrel @ 1000 ppm (T_2) recorded the highest total sugars (13.78%), followed by ethrel @ 500 ppm (T_1) with 13.29%, whereas untreated control fruits showed the lowest value (11.22%). Storage duration significantly affected total sugars, which increased from 11.82% on the 3rd day to 13.86% on the 6th day, followed by a gradual decline to 11.65% on the 12th day of storage. The interaction effect was also significant, with maximum total sugars (15.91%) observed in fruits treated with ethrel @ 1000 ppm on the 9th day, while minimum total sugars (9.39%) were recorded in CaC_2 treated fruits on the 12th day of storage. Ethrel @ 1000 ppm maintained comparatively higher sugar content even at the end of storage. The initial increase in total sugars during the process of ripening due to enzymatic hydrolysis of starch and polysaccharides into simple sugars, whereas the following decline was associated with utilization of sugars during respiration and senescence (Yu *et al.*, 2022). Ethrel treated fruits showed higher sugar accumulation due to enhanced ethylene mediated activation of ripening enzymes and improved conversion of reserve carbohydrates into soluble sugars (Tipu and Sherif, 2024). Moderate ethrel concentrations maintained higher sugar levels for longer duration, indicating balanced ripening and delayed metabolic depletion (Wardani *et al.*, 2026). Similar findings were reported by Tsomu *et al.* (2015), Kaur *et al.* (2017) while in mango by Palpandian *et al.* (2019).

Total Phenolic Content (mg GAE/100g FW)

Total phenolic content (TPC) of sapota cv. Cricket Ball was significantly influenced by ripening treatments, storage duration and their interaction (Fig. 3a). Among the treatments, untreated control fruits (T_7) recorded the highest phenolic content (135.10 mg GAE/100g FW), followed by ethrel @ 500 ppm (T_1), whereas CaC_2 treated fruits (T_6)

exhibited the lowest value (106.97 mg GAE/100g FW). Storage period significantly affected TPC, which declined progressively from 135.37 mg GAE/100g FW on the 3rd day to 105.57 mg GAE/100g FW on the 12th day of storage. The significant treatment $\times$ storage interaction indicated that the extent of change in TPC during storage varied among the ripening treatments. A gradual decline in TPC was observed throughout the ripening period, with CaC_2 treated fruits showed reduction compared to other treatments, while control fruits maintained comparatively higher phenolic content during storage. The decline in phenolic content during ripening and storage which may be attributed to enhanced oxidative degradation and polymerization of phenolic compounds during senescence (Zhang *et al.* 2023). Ethrel and calcium carbide treatments accelerated ripening, thereby increasing the activity of oxidative enzymes such as polyphenol oxidase and peroxidase responsible for rapid oxidation of phenolic constituents (Vidhya *et al.*, 2025; Chandar and Smith, 2019). Reduction in soluble tannins and conversion of phenolics into insoluble forms during ripening also contributed to lower detectable phenolic content (Shahidi and Hossain, 2023). Similar findings were reported by Torres Rodriguez *et al.* (2011) in sapote, Fernando *et al.* (2014) in banana, Rastegar *et al.* (2015) in sapota fruits.

Total Antioxidant Activity (%)

Total antioxidant activity of sapota cv. Cricket Ball was significantly influenced by ripening treatments, storage duration and their interaction (Fig. 3b). Among the treatments, untreated control fruits (T_7) recorded the maximum antioxidant activity (43.14%), followed by ethrel @ 500 ppm (T_1) with 37.19%, whereas CaC_2 treated fruits (T_6) exhibited the minimum value (20.85%). Storage duration significantly affected antioxidant activity, which declined progressively from 48.27% on the 3rd day to 18.95% on the 12th day of storage. The interaction effect was also significant, with maximum antioxidant activity (61.28%) observed in control fruits on the 3rd day, while the minimum value (12.09%) was recorded in CaC_2 treated fruits on the 12th day of storage. A continuous reduction in antioxidant activity was observed throughout

ripening, with CaC_2 treated fruits showing a sharper decline compared to other treatments. The reduction in antioxidant activity during ripening and storage attributed to degradation and oxidation of antioxidant compounds such as ascorbic acid, phenolics, carotenoids and other bioactive metabolites during senescence (de Souza *et al.*, 2021). Accelerated ripening induced by ethrel and calcium carbide enhanced oxidative metabolism and increased utilization of antioxidant constituents, leading to rapid decline in radical scavenging capacity (Vidhya *et al.*, 2025; Singh *et al.*, 2013). Lower antioxidant retention in CaC_2 fruits due to stress induced decline and enhanced oxidative enzyme activity (Mode *et al.*, 2026).

Shelf-life

Shelf-life of sapota cv. Cricket Ball was significantly influenced by different ripening treatments (Plates 1a & 1b). Among the treatments, fruits treated with ethrel @ 500 ppm (T_1) recorded the maximum shelf-life (10.50 days), which was statistically at par with ethrel @ 1000 ppm (T_2) exhibiting 10.33 days. However, CaC_2 treated fruits (T_6) showed the minimum shelf-life (6.50 days), differing




Fig. 3: Effect of post-harvest ethrel treatments and storage duration on a. total phenolic content (TPC) b. total antioxidant activity of sapota cv. Cricket Ball during ripening




Plate 1(a): Visual condition of sapota fruits at different storage intervals as affected by different treatments



Plate 1(b): Visual condition of sapota fruits at different storage intervals as affected by different treatments

significantly from all other treatments. Shelf life decreased progressively with increasing concentrations of ethrel (Ran *et al.*, 2022). The reduction in shelf life at higher ethrel concentrations because of accelerated ethylene mediated ripening, increased respiration rate, rapid moisture loss and enhanced metabolic degradation of fruit tissues (Zhang *et al.*, 2017). Excessive ethylene exposure hastened softening, senescence and deterioration of quality parameters such as firmness, sugars and acidity, thereby shortening marketable life (Nagaraja *et al.*, 2026). However, lower concentrations of ethrel maintained a balanced ripening process, ensuring desirable fruit quality with comparatively extended shelf-life. Untreated fruits retained firmness for longer duration but failed to attain acceptable ripening and marketable quality during storage. Although ethrel @ 500 ppm recorded the highest shelf-life, ethrel @ 1000 ppm was statistically at par and provided comparatively higher TSS and total sugar accumulation with acceptable fruit quality. Calcium carbide treated fruits exhibited the minimum shelf life due to rapid acetylene persuaded ripening, excessive tissue breakdown and faster senescence (Vidhya *et al.*, 2025).

CONCLUSION

The present study confirmed that postharvest application of ethrel significantly influenced the ripening behaviour, physicochemical quality, antioxidant properties and shelf-life of sapota cv. Cricket Ball under ambient storage conditions. Ethrel treated fruits exhibited faster and more uniform ripening compared to untreated fruits, but calcium carbide treatment resulted in rapid deterioration and poor storage quality. Among the treatments, ethrel @ 1000 ppm (T_2) was found to be the most effective, as it produced fruits with superior total soluble solids, higher total sugar, desirable ripening characteristics and better eating quality. However, the maximum shelf-life was recorded with ethrel @ 500 ppm (T_1) but it was statistically at par with T_2 . Additionally, T_2 maintained acceptable antioxidant properties and overall fruit quality throughout the storage period. Therefore, post-harvest application of ethrel @ 1000 ppm is recommended for achieving

uniform ripening, improved fruit quality and satisfactory shelf life in sapota fruits under ambient conditions.

REFERENCES

- Adeyemi, M. M., Bawa, M. H. and Muktar, B. (2018). Evaluation of the effect of calcium carbide on induced ripening of banana, pawpaw and mango cultivated within Kaduna metropolis, Nigeria. *Journal of Chemical Society of Nigeria*, 43(2).
- Asrey, R., Sharma, S., Barman, K., Prajapati, U., Negi, N. and Meena, N. K. (2023). Biological and postharvest interventions to manage the ethylene in fruit: A review. *Sustainable Food Technology*, 1(6): 803-826.
- Baidya, B. K., Mahato, A., Pattnaik, R. K. and Sethy, P. (2020). A review of physiological and biochemical changes related to ripening along with post-harvest handling and treatments of sapota. *Journal of Pharmacognosy and Phytochemistry*, 9(4): 2030-2035.
- Baskar, M., Hemalatha, G. and Muneeshwari, P. (2020). Traditional and medicinal importance of sapota – Review. *International Journal of Current Microbiology and Applied Sciences*, 9(1): 1711-1717.
- Bons, H. K., Boora, R. S. and Rattanpal, H. S. (2016). Studies on effect of various levels of ethephon on ripening of sapodilla fruits (*Manilkara achras* (Mill.) Fosberg). *International Journal of Agriculture, Environment and Biotechnology*, 9(4): 563-567.
- Chandar, M. M. and Smith, A. A. (2019). Artificial ripening of banana – An overview. *Research Journal of Pharmacy and Technology*, 12(10): 5101-5106.
- de Souza, A. V., de Mello, J. M., da Silva Favaro, V. F., dos Santos, T. G. F., dos Santos, G. P., de Lucca Sartori, D. and Ferrari Putti, F. (2021). Metabolism of bioactive compounds and antioxidant activity in bananas during

- ripening. *Journal of Food Processing and Preservation*, 45(11): e15959.
- Deshi, V. V., Siddiqui, M. W., Homa, F., Lata, D. and Singh, D. R. (2024). CaC₂-induced ripening: Unveiling the bitter truth behind sweet fruit. *Food Chemistry*, 455: 140097.
- Fernando, H. R. P., Srilaong, V., Pongprasert, N., Boonyaritthongchai, P. and Jitareerat, P. (2014). Changes in antioxidant properties and chemical composition during ripening in banana variety 'Hom Thong' (AAA group) and 'Khai' (AA group). *International Food Research Journal*, 21(2): 749-754.
- Ghimire, R., Yadav, P. K., Shrestha, A. K., Devkota, A. R. and Khanal, S. (2021). Ripening regulation of banana cv. Malbhog using different ripening inducers. *Turkish Journal of Agriculture-Food Science and Technology*, 9(12): 2113-2123.
- Kaur, J., Singh, A., Singh, B. and Sharma, S. (2020). Sapota. In: *Antioxidants in Fruits: Properties and Health Benefits*. Springer, Singapore, Pp: 181-199.
- Liu, H., He, J., Lyu, L., Wu, W., Tian, J., Wu, Y. and Li, W. (2026). Research progress on the mechanism of fruit ripening and softening. *Russian Journal of Plant Physiology*, 73(3): 110.
- Madani, B., Mirshekari, A., Yahia, E. and Golding, J. B. (2018). Sapota (*Manilkara achras* Forb.) factors influencing fresh and processed fruit quality. *Horticultural Reviews*, 45: 105-142.
- Meiyarasan, S., Sivaprasad, M., Sreedhar, D. and Kumar Reddy, Y. S. (2025). Quality enhancement and shelf-life extension in mango (*Mangifera indica* L.) cv. Banganapalli through plant growth regulators, chemical sprays and pruning. *Plant Archives*, 25(2).
- Menezes, J. and Athmaselvi, K. A. (2016). Study on effect of polysaccharide based edible coating on the shelf life of sapota fruits stored at room temperature. *Current Nutrition and Food Science*, 12(4): 272-278.
- MoA&FW. (2025). Final estimate: Area and production of horticulture crops. Department of Agriculture and Farmers' Welfare, Ministry of Agriculture and Farmers' Welfare, Government of India (Released on 07.02.2025).
- Mode, A. M., Magaji, U. F., Babatunde, M. S., Dandare, A., Dabai, I. A. and Dandare, S. U. (2026). Effects of chemical ripening with calcium carbide on nutritional composition, antioxidant capacity and heavy metal accumulation in climacteric fruits. *International Journal of Food Science*, 2026(1): 7226409.
- Muthal, T., Harini, R., Sunil, C. K. and Chidanand, D. V. (2019). Effect of artificial ripening techniques on physico-chemical properties of banana (*Musa* sp.) variety Poovan. *International Journal of Pure and Applied Bioscience*, 7(2): 482-489.
- Nagaraja, S. K., Kumar, P., R. K., Nabi, S. U., Mir, J. I., Verma, M. K. and Mansoor, S. (2026). A comprehensive review of biochemical insights and advanced packaging technologies for shelf-life enhancement of temperate fruits. *Biosensors*, 16(2): 94.
- Oladipupo, A., Akinleye, M. and Coker, H. (2022). The use of calcium carbide in fruit ripening: Health risks and arsenic index as a quantitative marker for calcium carbide residue. *Progress in Chemical and Biochemical Research*, 5(2): 125-132.
- Orisa, C. A. and Usoroh, C. I. (2020). Effects of accelerated ripening agents on the fruit quality of pawpaw (*Carica papaya*). *ARC Journal of Nutrition and Growth*, 6(2): 42-49.
- Padmaja, N., Bosco, S. J. D. and Rao, J. S. (2015). Physico chemical analysis of sapota (*Manilkara zapota*) coated by edible aloe vera gel. *International Journal of Applied Sciences and Biotechnology*, 3(1): 20-25.
- Palpandian, P., Shanmugam, H., Rani, E. A. and Prabu, G. T. V. (2019). Determination of fruit quality of calcium carbide induced ripening in mango (*Mangifera indica* L. cv. Alphonso) by physiological, biochemical, bio-enzymatic and elemental composition analysis (EDX). *Indian Journal of*

- Biochemistry and Biophysics*, 56: 205-213.
- Pareek, S. (2016). Ripening physiology: An overview. In: *Postharvest Ripening Physiology of Crops*. CRC Press, Pp: 1–48.
- Ran, Y., Zheng, Y., Li, X., Jia, X., Duan, L., Du, M. and Jiang, Y. (2022). Combination of ethrel and antiseptic medium flow of micro-circulating improves shelf-life quality of kiwifruit after long-term controlled atmosphere storage. *Journal of Food Processing and Preservation*, 46(12): e17243.
- Rana, S. S., Pradhan, R. C. and Mishra, S. (2021). Physicochemical and physiological changes during storage. In: *Packaging and Storage of Fruits and Vegetables*. Apple Academic Press, Pp: 1–24.
- Rastegar, S. (2015). Physical, biochemical and mineral evaluation of sapota fruits during growth, development and ripening. *Journal of Fruit and Ornamental Plant Research*, 23(1): 14-19.
- Rekha, A. (2024). Sapota. In: *Fruit and Nut Crops*. Springer Nature, Singapore, Pp: 491–504.
- Shahidi, F. and Hossain, A. (2023). Importance of insoluble-bound phenolics to the antioxidant potential is dictated by source material. *Antioxidants*, 12(1): 203.
- Singh, J., Bal, J. S., Singh, S. and Mirza, A. (2018). Assessment of chemicals and growth regulators on fruit ripening and quality: A review. *Journal of Pharmacognosy and Phytochemistry*, 7(2): 1215-1222.
- Singh, Z., Khan, A. S., Zhu, S. and Payne, A. D. (2013). Nitric oxide in the regulation of fruit ripening: Challenges and thrusts. *Stewart Postharvest Review*, 9(4): 1-11.
- Srivastava, A. K., Kashyap, P., Meena, V. S., Verma, N. and Singh, S. P. (2017). Sapota [*Manilkara achras* (Mill.) Fosberg syn. *Achras zapota* L.]. In: Ghosh, S. N., Singh, A. and Thakur, A. (eds). *Underutilized Fruit Crops: Importance and Cultivation*. Jaya Publishing House, New Delhi, Pp: 1159–1194.
- Tipu, M. M. and Sherif, S. M. (2024). Ethylene and its crosstalk with hormonal pathways in fruit ripening: Mechanisms, modulation and commercial exploitation. *Frontiers in Plant Science*, 15: 1475496.
- Torres-Rodríguez, A., Salinas-Moreno, Y., Valle-Guadarrama, S. and Alia-Tejagal, I. (2011). Soluble phenols and antioxidant activity in mamey sapote (*Pouteria sapota*) fruits in postharvest. *Food Research International*, 44(7): 1956-1961.
- Tsomu, T., Patel, H. C., Thakkar, R. M., Ajang, M. and Vasara, R. P. (2015). Response of post-harvest treatments of chemicals and plant growth regulators on biochemical characteristics of sapota fruit cv. Kalipatti. *The Bioscan*, 10(4): 1867-1870.
- Vidhya, D., Mahanti, N. K., Dakho, B., Kumar, A., Chaubey, S. and Chhetri, K. B. (2025). Calcium carbide (CaC₂) ripening in fruits: Health risks, non-destructive detection, quality control and regulatory frameworks. *Comprehensive Reviews in Food Science and Food Safety*, 24(2): e70140.
- Vidhya, M., Pandiarajan, T., Pandiselvam, R., Amirtham, D., Balakrishnan, M., Haseena, R. and Nishashree, R. (2017). Effect of ethylene concentration and exposure time on physico-chemical quality and colour value of sapota fruit (*Manilkara zapota*). *Asian Journal of Chemistry*, 29(5): 1045-1050.
- Wardani, N. P. Z. K., Suketi, K. and Arif, A. B. (2026). Ethrel-induced enhancement of sugar accumulation and postharvest quality in early-harvested cantaloupe melons. *The Scientific World Journal*, 2026(1): 9929514.
- Xia, Y., Wu, D. T., Ali, M., Liu, Y., Zhuang, Q. G., Wadood, S. A. and Gan, R. Y. (2024). Innovative postharvest strategies for maintaining the quality of kiwifruit during storage: An updated review. *Food Frontiers*, 5(5): 1933-1950.
- Yu, J., Tseng, Y., Pham, K., Liu, M. and Beckles, D. M. (2022). Starch and sugars as determinants of postharvest shelf life and quality: Some new and surprising roles. *Current Opinion in Biotechnology*, 78: 102844.

- Zhang, J., Cheng, D., Wang, B., Khan, I. and Ni, Y. (2017). Ethylene control technologies in extending postharvest shelf life of climacteric fruit. *Journal of Agricultural and Food Chemistry*, 65(34): 7308-7319.
- Zhang, Y. J., Huang, Q., Li, A. R., Gan, Z. Y., Zeng, J. K., Kai, W. B. and Chen, J. Y. (2023). Apple polyphenols delay postharvest

senescence and quality deterioration of 'Jinshayou' pummelo fruit during storage. *Frontiers in Plant Science*, 13: 1117106.

Received: June 30, 2026

Accepted: August 12, 2026